arXiv:math/9904082v1 [math.QA] 16 Apr 1999

FACE ALGEBRAS AND UNITARITY OF SU(N)$_{\mathrm{L}}$-TQFT

TAKAHIRO HAYASHI

Abstract. Using face algebras (i.e. algebras of L-operators of IRF models), we construct modular tensor categories with positive definite inner product, whose fusion rules and S-matrices are the same as (or slightly different from) those obtained by $U_q(\mathfrak{sl}_{\mathfrak{N}})$ at roots of unity. Also we obtain state-sums of ABF models on framed links which give quantum $SU(2)$-invariants of corresponding 3-manifolds.

1. INTRODUCTION

As is well known, quantum groups have their origin in the theory of quantum inverse scattering method. More specifically, they first appeared as so-called algebras of L-operators of lattice models (of vertex type). For example, the simplest quantum group $U_q(\mathfrak{sl}(2))$ can be viewed as the algebra of L-operators of 6-vertex model without spectral parameter. It seems that it is worth trying to study algebras of L-operators independently from the framework of Drinfeld-Jimbo algebra.

By investigating algebraic structure of lattice models of face type, we found a new class of quantum groups, which is called the class of *face algebras* (cf. [14]-[22] and also [3, 24, 36]). It contains all bialgebras as a subclass. Moreover, as well as bialgebras, face algebras produce monoidal categories as their (co-)module categories.

In this paper, we give a detailed study of face algebras $\mathfrak{S}(A_{N-1};t)_\epsilon$, which are obtained as algebras of L-operators of RSOS models of type A_{N-1} ($N \geq 2$) (cf. [23]), where $\epsilon = \pm 1$ and t denotes a primitive $2(N+L)$-th root of unity with $L \geq 1$. We also give two applications of $\mathfrak{S}(A_{N-1};t)_\epsilon$ to 3-dimensional topological quantum field theory (TQFT) and corresponding quantum invariants of 3-manifolds.

We show that the algebra $\mathfrak{S}(A_{N-1};t)_\epsilon$ is finite-dimensional cosemisimple and that its dual is a C^*-algebra for a suitable t. Also, we classify irreducible comodules of $\mathfrak{S}(A_{N-1};t)_\epsilon$ and determine their dimensions. Moreover, we construct various structures on $\mathfrak{S}(A_{N-1};t)_\epsilon$, such as the antipode, the braiding and the ribbon functional. The algebra $\mathfrak{S}(A_{N-1};t)_\epsilon$ is constructed as a quotient of the face version $\mathfrak{A}(w_{N,t,\epsilon})$ of FRT construction modulo one additional relation "det = 1", where $w_{N,t,\epsilon}$ is the Boltzmann weight of RSOS models of type A_{N-1} without the spectral parameter and det denotes the "(quantum) determinant" of $\mathfrak{A}(w_{N,t,\epsilon})$. Since the representation theory of $\mathfrak{A}(w_{N,t,\epsilon})$ is relatively easily established using a result on Iwahori-Hecke algebras due to H. Wenzl, the core of our work is to study the properties of the element det or, corresponding "exterior" algebra.

Next, we explain the unitarity of 3-dimensional TQFT briefly. Roughly speaking, a 3-dimensional TQFT is a map which assigns to each 3-cobordism $(M, \partial_- M, \partial_+ M)$, a linear map $\tau(M)\colon \mathcal{T}(\partial_-\mathcal{M}) \to \mathcal{T}(\partial_+\mathcal{M})$. Here, by a 3-dimensional cobordism, we mean a compact 3-dimensional manifold M whose boundary is a disjoint union of two closed surfaces $\partial_- M, \partial_+ M$. A 3-dimensional TQFT is called unitary if $\mathcal{T}(\partial_\pm\mathcal{M})$ are (finite-dimensional) Hilbert spaces and $\tau(-M) = \tau(M)^*$ for each 3-cobordism

---

Date:
Graduate School of Mathematics, Nagoya University,
Furo-cho, Chikusa-ku, Nagoya 464, Japan.

$(M, \partial_- M, \partial_+ M)$. It is established in [40] that to obtain a (unitary) 3-dimensional MTC, it suffices to construct a (unitary) modular tensor category (MTC) (i.e. a braided category which satisfies certain additional properties). The most important examples of MTC are constructed as semisimple quotients $\mathcal{C}(\mathfrak{g}, \mathfrak{q})$ of some module categories of Drinfeld-Jimbo algebra $U_q(\mathfrak{g})$ at roots q of unity (cf. [34, 1, 8, 28], and see also [7, 41, 42] for other construction of MTC's). For a suitable q, it also is expected that $\mathcal{C}(\mathfrak{g}, \mathfrak{q})$ is a unitary MTC. However, it seems that it is not easy to verify it directly, since $U_q(\mathfrak{g})$ is non-semisimple and cannot have a C^*-algebra structure (cf. A. Kirillov, Jr, [28] and V. Tuaev and H. Wenzl [42]).

The first application of $\mathfrak{S}(A_{N-1}; t)_\epsilon$ is to show that the category $\mathcal{C}_{\mathfrak{S}}(\mathcal{A}_{\mathcal{N}-\infty}, \sqcup)_\epsilon$ of all finite-dimensional right $\mathfrak{S}(A_{N-1}; t)_\epsilon$-comodules is a MTC, and that (for a suitable t), the category $\mathcal{C}^{\sqcap}_{\mathfrak{S}}(\mathcal{A}_{\mathcal{N}-\infty}, \sqcup)_\epsilon$ of all finite-dimensional unitary $\mathfrak{S}(A_{N-1}; t)_\epsilon$-comodules is a unitary MTC, whose fusion rules agree with those of $\mathcal{C}(\mathfrak{sl}_{\mathfrak{N}}, \mathfrak{q})$. Here "unitary" comodule means a comodule with inner product which satisfies some conditions. The quantum dimensions and S-matrices of $\mathcal{C}_{\mathfrak{S}}(\mathcal{A}_{\mathcal{N}-\infty}, \sqcup)_\epsilon$ are the same as (or slightly different from) those of $\mathcal{C}(\mathfrak{sl}_{\mathfrak{N}}, \mathfrak{q})$ (according to the choice of ϵ and another sign parameter ι when N is even). Although we use $U_q(\mathfrak{sl}_{\mathfrak{N}})$ to obtain some combinatorial formulas, the essential part of our theory is independent from $U_q(\mathfrak{sl}_{\mathfrak{N}})$. Hence the equivalence of $\mathcal{C}(\mathfrak{sl}_{\mathfrak{N}}, \mathfrak{q})$ and $\mathcal{C}_{\mathfrak{S}}(\mathcal{A}_{\mathcal{N}-\infty}, \sqcup)_\epsilon$ is left as an open problem. However, by the result of Kazhdan and Wenzl [27], these two categories are equivalent up to a "twist."

Unlike the module category of $U_q(\mathfrak{g})$, the category $\mathcal{C}_{\mathfrak{S}}(\mathcal{A}_{\mathcal{N}-\infty}, \sqcup)_\epsilon$ itself is semisimple. Moreover, it has a apparent similarity to the spaces of the conformal blocks of Wess-Zumino-Witten (WZW) models. We hope that there exists a direct connection between $\mathfrak{S}(A_{N-1}; t)_\epsilon$ and WZW models, similarly to Drinfeld [6].

The second application is to give an explicit description of the quantum $SU(2)$-invariant $\tau(M)$ of closed 3-manifolds M, which is associated with $\mathcal{C}_{\mathfrak{S}}(\mathcal{A}_{\infty}, \sqcup)_\epsilon$. More precisely, we express $\tau(M)$ as a state sum on each generic link diagram D which represents M. It gives a direct connection between the invariant and ABF model.

The paper is organized as follows. We start in Sect. 2-4, by recalling elementary properties of face algebras $\mathfrak{H}$, various structures on $\mathfrak{H}$ and their relations to the comodule category of $\mathfrak{H}$. In Sect. 5, we recall the notion of star-triangular (Yang-Baxter) face models and flat face models. The later is introduced in [18], and is a variant of Ocneanu's notion of flat biunitary connection in operator algebra. Flat face models play an crucial role to the determination of the representation theory of $\mathfrak{S}(A_{N-1}; t)_\epsilon$. In Sect. 6-7, we define the algebras $\mathfrak{S}(A_{N-1}; t)_\epsilon$ and state the main result on the representation theory of them. In Sect. 9-10, we construct several structures on $\mathfrak{S}(A_{N-1}; t)_\epsilon$, which we call the transpose, the costar structure, the antipode and the ribbon functional. Consequently, we see that $\mathcal{C}_{\mathfrak{S}}(\mathcal{A}_{\mathcal{N}-\infty}, \sqcup)_\epsilon$ (resp. $\mathcal{C}^{\sqcap}_{\mathfrak{S}}(\mathcal{A}_{\mathcal{N}-\infty}, \sqcup)_\epsilon$) is a (unitary) ribbon category. In Sect. 11, we prove that $\mathcal{C}_{\mathfrak{S}}(\mathcal{A}_{\mathcal{N}-\infty}, \sqcup)_\epsilon$ is a modular tensor category, by computing its S-matrix. Section 8 and Section 12 are devoted to some technical calculations on face analogues of the exterior algebras. In Sect. 13-14, we give an explicit description of $\mathcal{C}_{\mathfrak{S}}(\mathcal{A}_{\infty}, \sqcup)_\epsilon$ and give a state sum expression of the quantum $SU(2)$-invariant stated above.

After submitting the manuscript, one of the referee informed me the existence of the following two papers;

H. Wenzl, C^*-tensor categories from quantum groups, *J. Amer. Math. Soc.* **11** (1988), pp. 261-282.

C. Blanchet, Heck algebras, modular categories and 3-manifolds quantum invariants, preprint.

The former gives a proof of the unitarity of $\mathcal{C}(\mathfrak{g}, \mathfrak{q})$ for each $\mathfrak{g}$. The latter gives a construction of modular tensor categories via Iwahori-Hecke algebras at root of unity.

Throughout this paper, we use Sweedler's sigma notation for coalgebras, such as $(\Delta \otimes \mathrm{id})(\Delta(\mathrm{a})) = \sum_{(\mathrm{a})} \mathrm{a}_{(1)} \otimes \mathrm{a}_{(2)} \otimes \mathrm{a}_{(3)}$ (cf. [37]).

2. FACE ALGEBRAS

In this section, we give the definition of the face algebra and various structures on it. Let $\mathfrak{H}$ be an algebra over a field $\mathbb{K}$ equipped with a coalgebra structure $(\mathfrak{H}, \Delta, \varepsilon)$. Let $\mathcal{V}$ be a finite non-empty set and let e_λ and $\mathring{e}_\lambda$ $(\lambda \in \mathcal{V})$ be elements of $\mathfrak{H}$. We say that $(\mathfrak{H}, e_\lambda, \mathring{e}_\lambda)$ is a $\mathcal{V}$-*face algebra* if the following relations are satisfied:

$$\Delta(ab) = \Delta(a)\Delta(b), \tag{2.1}$$

$$e_\lambda e_\mu = \delta_{\lambda\mu} e_\lambda, \quad \mathring{e}_\lambda \mathring{e}_\mu = \delta_{\lambda\mu} \mathring{e}_\lambda, \quad e_\lambda \mathring{e}_\mu = \mathring{e}_\mu e_\lambda, \tag{2.2}$$

$$\sum_{\nu \in \mathcal{V}} e_\nu = 1 = \sum_{\nu \in \mathcal{V}} \mathring{e}_\nu, \tag{2.3}$$

$$\Delta(\mathring{e}_\lambda e_\mu) = \sum_{\nu \in \mathcal{V}} \mathring{e}_\lambda e_\nu \otimes \mathring{e}_\nu e_\mu, \quad \varepsilon(\mathring{e}_\lambda e_\mu) = \delta_{\lambda\mu}, \tag{2.4}$$

$$\varepsilon(ab) = \sum_{\nu \in \mathcal{V}} \varepsilon(a e_\nu) \varepsilon(\mathring{e}_\nu b) \tag{2.5}$$

for each $a, b \in \mathfrak{H}$ and $\lambda, \mu \in \mathcal{V}$. We call elements e_λ and $\mathring{e}_\lambda$ *face idempotents* of $\mathfrak{H}$. It is known that bialgebra is an equivalent notion of $\mathcal{V}$-face algebra with card($\mathcal{V}$) = . For a $\mathcal{V}$-face algebra $\mathfrak{H}$, we have the following formulas:

$$\varepsilon(\mathring{e}_\lambda a) = \varepsilon(e_\lambda a), \quad \varepsilon(a \mathring{e}_\lambda) = \varepsilon(a e_\lambda), \tag{2.6}$$

$$\sum_{(a)} a_{(1)} \varepsilon(e_\lambda a_{(2)} e_\mu) = e_\lambda a e_\mu, \tag{2.7}$$

$$\sum_{(a)} \varepsilon(e_\lambda a_{(1)} e_\mu) a_{(2)} = \mathring{e}_\lambda a \mathring{e}_\mu, \tag{2.8}$$

$$\Delta(a) = \sum_{\nu, \xi} \sum_{(a)} e_\nu a_{(1)} e_\xi \otimes \mathring{e}_\nu a_{(2)} \mathring{e}_\xi, \tag{2.9}$$

$$\sum_{(a)} e_\lambda a_{(1)} e_\mu \otimes a_{(2)} = \sum_{(a)} a_{(1)} \otimes \mathring{e}_\lambda a_{(2)} \mathring{e}_\mu, \tag{2.10}$$

$$\Delta(\mathring{e}_\lambda e_\mu a \mathring{e}_{\lambda'} e_{\mu'}) = \sum_{(a)} \mathring{e}_\lambda a_{(1)} \mathring{e}_{\lambda'} \otimes e_\mu a_{(2)} e_{\mu'} \tag{2.11}$$

for each $a \in \mathfrak{H}$ and $\lambda, \mu, \lambda', \mu' \in \mathcal{V}$.

Let $\mathcal{G}$ be a finite oriented graph with set of vertices $\mathcal{V} = \mathcal{G}^0$. For an edge $\mathbf{p}$, we denote by $\mathfrak{s}(\mathbf{p})$ and $\mathfrak{r}(\mathbf{p})$ its *source* (*start*) and its *range* (*end*) respectively. For each $m \geq 1$, we denote by $\mathcal{G}^m = \coprod_{\lambda, \mu \in \mathcal{V}} \mathcal{G}^m_{\lambda\mu}$ the set of *paths* of $\mathcal{G}$ of *length* m, that is, $\mathbf{p} \in \mathcal{G}^{\mathbf{m}}_{\lambda\mu}$ if $\mathbf{p}$ is a sequence $(\mathbf{p_1}, \ldots, \mathbf{p_m})$ of edges of $\mathcal{G}$ such that $\mathfrak{s}(\mathbf{p}) := \mathfrak{s}(\mathbf{p_1}) = \lambda$, $\mathfrak{r}(\mathbf{p_n}) = \mathfrak{s}(\mathbf{p_{n+1}})$ $(1 \leq n < m)$ and $\mathfrak{r}(\mathbf{p}) := \mathfrak{r}(\mathbf{p_m}) = \mu$. Let $\mathfrak{H}(\mathcal{G})$ be the linear span of the symbols $e\binom{\mathbf{p}}{\mathbf{q}}$ $(\mathbf{p}, \mathbf{q} \in \mathcal{G}^{\mathbf{m}}, \mathbf{m} \geq \mathbf{0})$. Then $\mathfrak{H}(\mathcal{G})$ becomes a $\mathcal{V}$-face algebra by setting

$$\mathring{e}_\lambda = \sum_{\mu \in \mathcal{V}} e\binom{\lambda}{\mu}, \quad e_\mu = \sum_{\lambda \in \mathcal{V}} e\binom{\lambda}{\mu}, \tag{2.12}$$

$$e\binom{\mathbf{p}}{\mathbf{q}} e\binom{\mathbf{r}}{\mathbf{s}} = \delta_{\mathfrak{r}(\mathbf{p})\mathfrak{s}(\mathbf{r})}\, \delta_{\mathfrak{r}(\mathbf{q})\mathfrak{s}(\mathbf{s})}\, e\binom{\mathbf{p} \cdot \mathbf{r}}{\mathbf{q} \cdot \mathbf{s}}, \tag{2.13}$$

$$\Delta\left(e\binom{\mathbf{p}}{\mathbf{q}}\right) = \sum_{\mathbf{t}\in\mathcal{G}^m} e\binom{\mathbf{p}}{\mathbf{t}} \otimes e\binom{\mathbf{t}}{\mathbf{q}}, \quad \varepsilon\left(e\binom{\mathbf{p}}{\mathbf{q}}\right) = \delta_{\mathbf{pq}} \tag{2.14}$$

for each $\mathbf{p}, \mathbf{q} \in \mathcal{G}^{\mathbf{m}}$ and $\mathbf{r}, \mathbf{s} \in \mathcal{G}^{\mathbf{n}}$ $(m, n \geq 0)$. Here for paths $\mathbf{p} = (\mathbf{p_1}, \dots, \mathbf{p_m})$ and $\mathbf{r} = (\mathbf{r_1}, \dots, \mathbf{r_n})$, we set $\mathbf{p}\cdot\mathbf{r} = (\mathbf{p_1}, \dots, \mathbf{p_m}, \mathbf{r_1}, \dots, \mathbf{r_n})$ if $\mathfrak{r}(\mathbf{p}) = \mathfrak{s}(\mathbf{r})$. We note that $\mathbb{K}\mathcal{G} = \bigoplus_{\mathbf{p}\in\mathcal{G}} \mathbb{K}\mathbf{p}$ $(m \geq 0)$ becomes a right $\mathfrak{H}(\mathcal{G})$-comodule via

$$\mathbf{q} \mapsto \sum_{\mathbf{p}\in\mathcal{G}} \mathbf{p} \otimes \mathbf{e}\binom{\mathbf{p}}{\mathbf{q}}. \tag{2.15}$$

Proposition 2.1 ([16])**.** *Every finitely generated* $\mathcal{V}$*-face algebra is isomorphic to a quotient of* $\mathfrak{H}(\mathcal{G})$ *for some* $\mathcal{G}$*.*

We say that a linear map $S : \mathfrak{H} \to \mathfrak{H}$ is an *antipode* of $\mathfrak{H}$, or $(\mathfrak{H}, S)$ is a *Hopf* $\mathcal{V}$*-face algebra* if

$$\sum_{(a)} S(a_{(1)})a_{(2)} = \sum_{\nu\in\mathcal{V}} \varepsilon(ae_\nu)e_\nu, \quad \sum_{(a)} a_{(1)}S(a_{(2)}) = \sum_{\nu\in\mathcal{V}} \varepsilon(e_\nu a)\mathring{e}_\nu, \tag{2.16}$$

$$\sum_{(a)} S(a_{(1)})a_{(2)}S(a_{(3)}) = S(a) \tag{2.17}$$

for each $a \in \mathfrak{H}$. An antipode of a $\mathcal{V}$-face algebra is an antialgebra-anticoalgebra map, which satisfies

$$S(\mathring{e}_\lambda e_\mu) = \mathring{e}_\mu e_\lambda \quad (\lambda, \mu \in \mathcal{V}). \tag{2.18}$$

The antipode of a $\mathcal{V}$-face algebra is unique if it exists.

Let $\mathfrak{H}$ be a $\mathcal{V}$-face algebra with product m and let $\mathcal{R}^+$ and $\mathcal{R}^-$ be elements of $(\mathfrak{H} \otimes \mathfrak{H})^*$. We say that $(\mathfrak{H}, \mathcal{R}^\pm)$ is a *coquasitriangular* (or *CQT*) $\mathcal{V}$-face algebra if the following relations are satisfied

$$\mathcal{R}^+ m^*(1) = \mathcal{R}^+, \quad m^*(1)\mathcal{R}^- = \mathcal{R}^-, \tag{2.19}$$

$$\mathcal{R}^-\mathcal{R}^+ = m^*(1), \quad \mathcal{R}^+\mathcal{R}^- = (m^{\mathrm{op}})^*(1), \tag{2.20}$$

$$\mathcal{R}^+ m^*(X)\mathcal{R}^- = (m^{\mathrm{op}})^*(X) \quad (X \in \mathfrak{H}^*), \tag{2.21}$$

$$(m\otimes\mathrm{id})^*(\mathcal{R}^+) = \mathcal{R}^+_{13}\mathcal{R}^+_{23}, \quad (\mathrm{id}\otimes m)^*(\mathcal{R}^+) = \mathcal{R}^+_{13}\mathcal{R}^+_{12}. \tag{2.22}$$

Here, as usual, for each $Z \in (\mathfrak{H}\otimes\mathfrak{H})^*$ and $\{i, j, k\} = \{1, 2, 3\}$, we define $Z_{ij} \in (\mathfrak{H}^{\otimes 3})^*$ by $Z_{ij}(a_1, a_2, a_3) = Z(a_i, a_j)\varepsilon(a_k)$ $(a_1, a_2, a_3 \in \mathfrak{H})$. We note, for example, that the second formula of (2.22) is equivalent to

$$\mathcal{R}^+(a,\, bc) = \sum_{(a)} \mathcal{R}^+(a_{(1)},\, c)\mathcal{R}^+(a_{(2)},\, b) \quad (a, b, c \in \mathfrak{H}). \tag{2.23}$$

It is known that $\mathcal{R}^\pm$ satisfies

$$(m\otimes\mathrm{id})^*(\mathcal{R}^-) = \mathcal{R}^-_{23}\mathcal{R}^-_{13}, \quad (\mathrm{id}\otimes m)^*(\mathcal{R}^-) = \mathcal{R}^-_{12}\mathcal{R}^-_{13}, \tag{2.24}$$

$$\mathcal{R}^+(\mathring{e}_\lambda e_\mu a \mathring{e}_\nu e_\xi,\, b) = \mathcal{R}^+(a,\, \mathring{e}_\mu e_\xi b \mathring{e}_\lambda e_\nu), \tag{2.25}$$

$$\mathcal{R}^-(\mathring{e}_\lambda e_\mu a \mathring{e}_\nu e_\xi,\, b) = \mathcal{R}^-(a,\, \mathring{e}_\nu e_\lambda b \mathring{e}_\xi e_\mu), \tag{2.26}$$

$$\mathcal{R}^+(\mathring{e}_\lambda e_\mu,\, a) = \varepsilon(e_\mu a e_\lambda), \quad \mathcal{R}^+(a,\, \mathring{e}_\lambda e_\mu) = \varepsilon(e_\lambda a e_\mu), \tag{2.27}$$

$$\mathcal{R}^-(\mathring{e}_\lambda e_\mu,\, a) = \varepsilon(e_\lambda a e_\mu), \quad \mathcal{R}^-(a,\, \mathring{e}_\lambda e_\mu) = \varepsilon(e_\mu a e_\lambda) \tag{2.28}$$

for each $\lambda, \mu, \nu, \xi \in \mathcal{V}$, $a, b \in \mathfrak{H}$. If, in addition, $\mathfrak{H}$ has an antipode S, then

$$(S \otimes \mathrm{id})^*(\mathcal{R}^+) = \mathcal{R}^-, \quad (\mathrm{id} \otimes S)^*(\mathcal{R}^-) = \mathcal{R}^+. \tag{2.29}$$

For a CQT Hopf $\mathcal{V}$-face algebra $\mathfrak{H}$, we define its *Drinfeld functionals* $\mathcal{U}_i \in \mathfrak{H}^*$ $(i = 1, 2)$ via

$$\mathcal{U}_1(a) = \sum_{(a)} \mathcal{R}^+(a_{(2)}, S(a_{(1)})), \quad \mathcal{U}_2(a) = \sum_{(a)} \mathcal{R}^-(S(a_{(1)}), a_{(2)}) \quad (a \in \mathfrak{H}). \tag{2.30}$$

The Drinfeld functionals are invertible as elements of the dual algebra $\mathfrak{H}^*$ and satisfy the following relations [20]:

$$\mathcal{U}_1^{-1}(a) = \sum_{(a)} \mathcal{R}^-(S(a_{(2)}), a_{(1)}), \quad \mathcal{U}_2^{-1}(a) = \sum_{(a)} \mathcal{R}^+(a_{(1)}, S(a_{(2)})), \tag{2.31}$$

$$\mathcal{U}_i^{\pm 1}(\overset{\circ}{e}_\lambda e_\mu) = \delta_{\lambda\mu}, \quad \mathcal{U}_i^{\pm}(\overset{\circ}{e}_\lambda a \overset{\circ}{e}_\mu) = \mathcal{U}_i^{\pm}(e_\lambda a e_\mu), \tag{2.32}$$

$$\begin{aligned} m^*(\mathcal{U}_1) &= \mathcal{R}^-\mathcal{R}^-_{21}(\mathcal{U}_1 \otimes \mathcal{U}_1) = (\mathcal{U}_1 \otimes \mathcal{U}_1)\mathcal{R}^-\mathcal{R}^-_{21}, \\ m^*(\mathcal{U}_2) &= \mathcal{R}^+_{21}\mathcal{R}^+(\mathcal{U}_2 \otimes \mathcal{U}_2) = (\mathcal{U}_2 \otimes \mathcal{U}_2)\mathcal{R}^+_{21}\mathcal{R}^+, \end{aligned} \tag{2.33}$$

$$\mathcal{U}_i X \mathcal{U}_i^{-1} = (S^2)^*(X), \tag{2.34}$$

$$S^*(\mathcal{U}_1^{\pm 1}) = \mathcal{U}_2^{\mp 1}, \quad S^*(\mathcal{U}_2^{\pm 1}) = \mathcal{U}_1^{\mp 1} \tag{2.35}$$

for each $i = 1, 2$, $a \in \mathfrak{H}$, $X \in \mathfrak{H}^*$ and $\lambda, \mu \in \mathcal{V}$. In particular, S is bijective and $\mathcal{U}_1\mathcal{U}_2^{-1}$ is a central element of $\mathfrak{H}^*$. Let $(\mathfrak{H}, \mathcal{R}^{\pm})$ be a CQT Hopf $\mathcal{V}$-face algebra and $\mathcal{V}$ an invertible central element of $\mathfrak{H}^*$. We say that $\mathcal{V}$ is a *ribbon functional* of $\mathfrak{H}$, or $(\mathfrak{H}, \mathcal{V})$ is a *coribbon Hopf* $\mathcal{V}$*-face algebra* if

$$m^*(\mathcal{V}) = \mathcal{R}^-\mathcal{R}^-_{21}(\mathcal{V} \otimes \mathcal{V}), \tag{2.36}$$

$$S^*(\mathcal{V}) = \mathcal{V}. \tag{2.37}$$

Let g (resp. $\mathcal{G}$) be an element of (resp. a linear functional on) a $\mathcal{V}$-face algebra $\mathfrak{H}$. We say that g (resp. $\mathcal{G}$) is *group-like* if the following relations (2.38)-(2.39) (resp. (2.40)-(2.41)) are satisfied:

$$\Delta(g) = \sum_{\nu \in \mathcal{V}} g e_\nu \otimes g \overset{\circ}{e}_\nu, \tag{2.38}$$

$$g \overset{\circ}{e}_\lambda e_\mu = \overset{\circ}{e}_\lambda e_\mu g, \quad \varepsilon(g \overset{\circ}{e}_\lambda e_\mu) = \delta_{\lambda\mu}, \tag{2.39}$$

$$\mathcal{G}(ab) = \sum_{\nu \in \mathcal{V}} \mathcal{G}(a e_\nu)\mathcal{G}(\overset{\circ}{e}_\nu b), \tag{2.40}$$

$$\mathcal{G}(\overset{\circ}{e}_\lambda a \overset{\circ}{e}_\mu) = \mathcal{G}(e_\lambda a e_\mu), \quad \mathcal{G}(\overset{\circ}{e}_\lambda e_\mu) = \delta_{\lambda\mu} \tag{2.41}$$

for each $a, b \in \mathfrak{H}$ and $\lambda, \mu \in \mathcal{V}$. If $\mathfrak{H}$ has an antipode S, then every group-like element g and group-like functional $\mathcal{G}$ are invertible and satisfy

$$S(g) = g^{-1}, \quad S^*(\mathcal{G}) = \mathcal{G}^{-1}. \tag{2.42}$$

We denote by GLE($\mathfrak{H}$) the set of all group-like elements of $\mathfrak{H}$.

Proposition 2.2 ([21])**.** *Let $\mathfrak{H}$ be a CQT Hopf $\mathcal{V}$-face algebra and $\mathcal{V}$ an invertible element of $\mathfrak{H}^*$. Then $(\mathfrak{H}, \mathcal{V})$ is a coribbon Hopf $\mathcal{V}$-face algebra if and only if $\mathcal{M} = \mathcal{U}_\infty \mathcal{V}^{-\infty}$ is group-like and satisfies the following relations*:

$$\mathcal{M} X \mathcal{M}^{-1} = (S^2)^*(X) \quad (X \in \mathfrak{H}^*), \tag{2.43}$$

$$\mathcal{M}^{\in} = \mathcal{U}_\infty \mathcal{U}_{\in}. \tag{2.44}$$

For a coribbon Hopf $\mathcal{V}$-face algebra $(\mathfrak{H}, \mathcal{V})$, we call $\mathcal{M} = \mathcal{U}_\infty \mathcal{V}^{-\infty}$ the *modified ribbon functional* of $\mathfrak{H}$ corresponding to $\mathcal{V}$.

Example 2.1. When $L = 1$, the algebra $\mathfrak{S}(A_{N-1}; t)_\epsilon$ is rather degenerate. Hence, we treat this case here separately from the case $L \geq 2$. (cf. [11]). Let $N \geq 2$ be an integer and $\mathcal{V}$ the cyclic group $\mathbb{Z}/\mathbb{N}\mathbb{Z}$. Let $t \in \mathbb{C}$ be a primitive $2(N+1)$-th root of unity and ϵ either 1 or -1. We define the $\mathcal{V}$-face algebra $\mathfrak{S}(A_{N-1}; t)_\epsilon$ to be the $\mathbb{C}$-linear span of the symbols $e^i_j(m)$ $(i, j, m \in \mathcal{V})$ equipped with the structure of $\mathcal{V}$-face algebra given by

$$e^i_j[p] e^k_l[q] = \delta_{i+p,k} \delta_{j+q,l} e^i_j[p+q] \quad (p, q \in \mathbb{Z}_{\geq \nvdash}), \tag{2.45}$$

$$\Delta(e^i_k(m)) = \sum_j e^i_j(m) \otimes e^j_k(m), \quad \varepsilon(e^i_j(m)) = \delta_{ij}, \tag{2.46}$$

$$\mathring{e}_i = \sum_j e^i_j(0), \quad e_j = \sum_i e^i_j(0). \tag{2.47}$$

Here, in (2.45), we set

$$e^i_j[qN + r] = (-\epsilon)^{(i-j)q(N-1)} e^i_j(r + N\mathbb{Z}) \tag{2.48}$$

for each $q \in \mathbb{Z}$ and $0 \leq r < N$. The algebra $\mathfrak{S}(A_{N-1}; t)_\epsilon$ becomes a coribbon Hopf $\mathcal{V}$-face algebra via

$$S(e^i_j[p]) = e^{j+p}_{i+p}[-p], \tag{2.49}$$

$$\mathcal{R}^+(e^i_j[p], e^k_l[q]) = \delta_{i,k+q} \delta_{jk} \delta_{i+p,l+q} \delta_{j+p,l} (-\zeta t)^{-pq}, \tag{2.50}$$

$$\mathcal{V}_\iota(\rceil^{\rangle}_{|}[_{\surd}]) = \delta_{\rangle|} \iota \surd (-\zeta \sqcup)^{\surd^{\in}}, \tag{2.51}$$

where $i, j, k, l \in \mathcal{V}$, $p, q \in \mathbb{Z}_{\geq \nvdash}$, ζ denotes a solution of $\zeta^N = \epsilon^{N-1} t$, $\iota = \pm 1$ if $N \in 2\mathbb{Z}$ and $\iota = 1$ if $N \in 1 + 2\mathbb{Z}$.

3. COMODULES OF FACE ALGEBRAS

In this section, we recall categorical properties of comodules of face algebras (cf. [16]). We refer the readers to [26] for the terminologies on monoidal (or tensor) categories.

Let M be a right comodule of a $\mathcal{V}$-face algebra $\mathfrak{H}$. We define its *face space decomposition* $M = \bigoplus_{\lambda, \mu \in \mathcal{V}} M(\lambda, \mu)$ by

$$M(\lambda, \mu) = \Big\{ \sum_{(u)} u_{(0)} \varepsilon(e_\lambda u_{(1)} e_\mu) \Bigm| u \in M \Big\}. \tag{3.1}$$

Here we denote the coaction $M \to M \otimes \mathfrak{H}$ by $u \mapsto \sum_{(u)} u_{(0)} \otimes u_{(1)}$ $(u \in M)$. Let N be another $\mathfrak{H}$-comodule. We define the *truncated tensor product* $M \bar{\otimes} N$ to be the $\mathfrak{H}$-comodule given by

$$(M \bar{\otimes} N)(\lambda, \mu) = \bigoplus\nolimits_{\nu \in \mathcal{V}} M(\lambda, \nu) \otimes N(\nu, \mu), \tag{3.2}$$

$$u \otimes v \mapsto \sum_{(u),(v)} u_{(0)} \otimes v_{(0)} \otimes u_{(1)} v_{(1)} \qquad (u \in M(\lambda, \nu), v \in N(\nu, \mu), \lambda, \mu, \nu \in \mathcal{V}). \tag{3.3}$$

With this operation, the category $\mathbf{Com}^{\mathbf{f}}_{\mathfrak{H}}$ of all finite-dimensional right $\mathfrak{H}$-comodules becomes a monoidal (or tensor) category whose unit object $\mathbb{K}\mathcal{V}$ is given by

$$\mathbb{K}\mathcal{V} = \bigoplus_{\mu \in \mathcal{V}} \mathbb{K}\mu; \quad \mu \mapsto \sum_{\lambda \in \mathcal{V}} \lambda \otimes {}^\circ{}_{\lambda\mu} \quad (\mu \in \mathcal{V}). \tag{3.4}$$

If, in addition, $\mathfrak{H}$ is CQT, then $\mathbf{Com}^{\mathbf{f}}_{\mathfrak{H}}$ becomes a braided monoidal category with braiding $c_{MN}\colon M\bar{\otimes}N \cong N\bar{\otimes}M$ given by

$$c_{MN}(u\otimes v) = \sum_{(u),(v)} v_{(0)}\otimes u_{(0)}\mathcal{R}^{+}(u_{(1)}, v_{(1)})$$
$$(u\in M(\lambda,\nu), v\in N(\nu,\mu), \lambda,\mu,\nu\in\mathcal{V}). \tag{3.5}$$

Next, suppose that $\mathfrak{H}$ has a bijective antipode and M is finite-dimensional. Then there exists a unique right $\mathfrak{H}$-comodule $M^{\vee}$ such that whose underlying vector space is the dual of M and that the coaction satisfies

$$\sum_{(u)}\langle v, u_{(0)}\rangle S(u_{(1)}) = \sum_{(v)}\langle v_{(0)}, u\rangle v_{(1)} \quad (u\in M, v\in M^{\vee}). \tag{3.6}$$

As vector spaces, we have

$$M^{\vee}(\lambda,\mu)\cong M(\mu,\lambda)^{*} \quad (\lambda,\mu\in\mathcal{V}). \tag{3.7}$$

The comodule $M^{\vee}$ becomes a left dual object of M via maps

$$b_M\colon \mathbb{K}\mathcal{V}\to\mathcal{M}\bar{\otimes}\mathcal{M}^{\vee};\quad \lambda\mapsto\sum_{\mu\in\mathcal{V}}\sum_{\mathbf{p}\in\mathcal{M}_{\lambda\mu}}\mathbf{p}\otimes\mathbf{p}^{\vee} \quad (\lambda\in\mathcal{V}), \tag{3.8}$$

$$d_M\colon M^{\vee}\bar{\otimes}M\to\mathbb{K}\mathcal{V};\quad \mathbf{p}^{\vee}\otimes\mathbf{q}\mapsto\delta_{\mathbf{pq}}\,\mu \quad (\mathbf{p}\in\mathcal{M}_{\nu\lambda}, \mathbf{q}\in\mathcal{M}_{\nu\mu}), \tag{3.9}$$

where $\mathcal{M}_{\lambda\mu}$ denotes a basis of $M(\lambda,\mu)$ and $\{\mathbf{p}^{\vee}\,|\,\mathbf{p}\in\mathcal{M}_{\lambda\mu}\}$ its dual basis. By replacing S in (3.6) with S^{-1}, we obtain another $\mathfrak{H}$-comodule structure on M^{*}, which gives the right dual $M^{\wedge}$ of M. We note that the canonical linear isomorphism $I_{M^{\wedge}}\colon M^{\wedge}\to M^{\vee}$ satisfies

$$I_{M^{\wedge}}(Xu) = (S^{-2})^{*}(X)I_{M^{\wedge}}(u) \tag{3.10}$$

for each $X\in\mathfrak{H}^{*}$ and $u\in M^{\wedge}$. Here, as usual, we regard M as a left $\mathfrak{H}^{*}$-module via

$$Xu = \sum_{(u)} u_{(0)}\langle X, u_{(1)}\rangle \quad (u\in M, X\in\mathfrak{H}^{*}). \tag{3.11}$$

Finally, suppose that $\mathfrak{H}$ is a coribbon Hopf $\mathcal{V}$-face algebra. Then $\mathbf{Com}^{\mathbf{f}}_{\mathfrak{H}}$ becomes a ribbon category with twist $\theta_M\colon M\cong M$ given by

$$\theta_M(u) = \mathcal{V}^{-\infty}\sqcap \quad (\sqcap\in\mathscr{M}). \tag{3.12}$$

Lemma 3.1. *Let $\mathfrak{H}$ be a coribbon Hopf $\mathcal{V}$-face algebra with modified ribbon functional $\mathscr{M}$ such that its unit comodule $\mathbb{K}\mathcal{V}$ is absolutely irreducible. Then, the quantum trace of the ribbon category $\mathbf{Com}^{\mathbf{f}}_{\mathfrak{H}}$ is given by*

$$\mathrm{Tr_q(f)} = \frac{1}{\mathrm{card}(\mathcal{V})}\mathrm{Tr}(\mathscr{M}\{) \tag{3.13}$$
$$= \mathrm{Tr}\left((\mathscr{M}\{)|_{\mathscr{M}(\lambda,-)}\right) \tag{3.14}$$

for each $M\in\mathrm{ob}(\mathbf{Com}^{\mathbf{f}}_{\mathfrak{H}})$ and $f\in\mathrm{End}_{\mathfrak{H}}(\mathrm{M})$, where λ is an arbitrary element of $\mathcal{V}$ and $M(\lambda,-)=\bigoplus_{\mu}M(\lambda,\mu)$.

Proof. We define $u_M\colon M\to M^{\vee\vee}$ by the composition

$$M\cong M\bar{\otimes}\mathbb{K}\mathcal{V}\xrightarrow{\mathrm{id}\bar{\otimes}\mathrm{b}}\mathcal{M}\bar{\otimes}\mathcal{M}^{\vee}\bar{\otimes}\mathcal{M}^{\vee\vee}\xrightarrow{\bar{\otimes}\mathrm{id}}$$
$$M^{\vee}\bar{\otimes}M\bar{\otimes}M^{\vee\vee}\xrightarrow{d\bar{\otimes}\mathrm{id}}\mathbb{K}\mathcal{V}\bar{\otimes}\mathcal{M}^{\vee\vee}\cong\mathcal{M}^{\vee\vee}.$$

Then, as a consequence of the fact that $\mathbf{Com}^{\mathbf{f}}_{\mathfrak{H}}$ is a rigid braided monoidal category, we have

$$d_M\circ c_{MM^{\vee}} = d_{M^{\vee}}\circ(u_M\bar{\otimes}\mathrm{id_{M^{\vee}}}). \tag{3.15}$$

On the other hand, since $u_M(v) = I_M(\mathcal{U}_\infty\sqsubseteq)$ $(v \in M)$, we have

$$(u_M \circ \theta_M)(v) = I_M(\mathcal{M}\sqsubseteq) \quad (\sqsubseteq \in \mathcal{M}), \tag{3.16}$$

where I_M is as in (3.10). Hence, by the definition of $\mathrm{Tr_q}$, we obtain

$$\begin{aligned}\mathrm{Tr_q(f)}(\lambda) &= d_{M^\vee} \circ (u_M\theta_M f \bar{\otimes} \mathrm{id_{M^\vee}}) \circ \mathrm{b_M}(\lambda)\\ &= \sum_{\mu\in\mathcal{V}} \sum_{\mathbf{p}\in\mathcal{M}_{\lambda\mu}} \langle I_M(\mathcal{M}\{(\mathbf{p})), \mathbf{p}^\vee\rangle\, \lambda\\ &= \mathrm{Tr}\left((\mathcal{M}\{)|_{\mathcal{M}(\lambda,-)}\right)\lambda\end{aligned}$$

as required, where $\mathcal{M}_{\lambda\mu}$ and $\mathbf{p}^\vee$ are as in (3.8). □

We say that a right comodule M of a $\mathcal{V}$-face algebra $\mathfrak{H}$ is *group-like* if $\dim(M(\lambda,\mu)) = \delta_{\lambda\mu}$ for each $\lambda, \mu \in \mathcal{V}$. For each $g \in \mathrm{GLE}(\mathfrak{H})$, we can define a group-like comodule $\mathbb{K}\mathcal{V}\} = \bigoplus_{\mu\in\mathcal{V}} \mathbb{K}\mu\eth$ by the coaction $\mu g \mapsto \sum_\lambda \lambda g \otimes \mathring{e}_\lambda e_\mu g$. Conversely, we have the following.

Lemma 3.2 ([19]). *For each group-like comodule M and its basis $g_\mu \in M(\mu,\mu)$ $(\mu \in \mathcal{V})$, we obtain a group-like element g by $\mathring{e}_\lambda e_\mu g = g\binom{\lambda}{\mu}$ and $g_\mu \mapsto \sum_\lambda g_\lambda \otimes g\binom{\lambda}{\mu}$. Moreover, every group-like element is obtained in this manner.*

Let $\mathfrak{H}$ be a $\mathcal{V}$-face algebra and $\{L_\psi \mid \psi \in \mathbf{\Lambda}\}$ its finite-dimensional comodules such that $\mathfrak{H} \cong \bigoplus_{\psi\in\mathbf{\Lambda}} \mathrm{End(L_\psi)}^*$ as coalgebras. Let g be a central group-like element of $\mathfrak{H}$. We say that g is *simply reducible* if there exists a set $\overline{\mathbf{\Lambda}}$ and a bijection $\varphi\colon \overline{\mathbf{\Lambda}} \times \mathbb{Z}_{\geq\not{\vdash}} \cong \mathbf{\Lambda}$ such that $L_{\varphi(\lambda,n)} \cong \mathbb{K}\mathcal{V}\}^{\backslash}\bar{\otimes}\mathcal{L}_{\varphi(\lambda,)}$ for each $\lambda \in \overline{\mathbf{\Lambda}}$ and $n \geq 0$.

Lemma 3.3 ([18]). *Let $\mathfrak{H}$, g etc. be as above. Then,* (1) *the element g is not a zero-divisor of $\mathfrak{H}$.* (2) *The quotient $\overline{\mathfrak{H}} = \mathfrak{H}/(\mathfrak{g} - \imath)$ is isomorphic to $\bigoplus_{\lambda\in\overline{\mathbf{\Lambda}}} \mathrm{End(L_\lambda)}^*$ as coalgebras, where L_λ is $L_{\varphi(\lambda,0)}$ viewed as an $\overline{\mathfrak{H}}$-comodule. As $\overline{\mathfrak{H}}$-comodules, we have $L_{\varphi(\lambda,n)} \cong L_\lambda$ for each $\lambda \in \overline{\mathbf{\Lambda}}$ and n.*

4. COMPACT FACE ALGEBRAS

Let $\mathfrak{H}$ be a $\mathcal{V}$-face algebra over the complex number field $\mathbb{C}$ and $\times : \mathfrak{H} \to \mathfrak{H}$; $a \mapsto a^\times$ an antilinear map. We say that $(\mathfrak{H}, \times)$ is a *costar $\mathcal{V}$-face algebra* [17] if

$$(a^\times)^\times = a, \quad (ab)^\times = a^\times b^\times, \tag{4.1}$$

$$\Delta(a^\times) = \sum_{(a)} a^\times_{(2)} \otimes a^\times_{(1)}, \tag{4.2}$$

$$e^\times_\lambda = \mathring{e}_\lambda \tag{4.3}$$

for each $\lambda \in \mathcal{V}$ and $a, b \in \mathfrak{H}$. Let $[t_{pq}]_{p,q\in I}$ be a finite-size matrix whose entries are elements of $\mathfrak{H}$. Then $[t_{pq}]$ is called a *unitary matrix corepresentation* if $\Delta(t_{pq}) = \sum_r t_{pr} \otimes t_{rq}$, $\varepsilon(t_{pq}) = \delta_{pq}$ and $t^\times_{pq} = t_{qp}$. A costar $\mathcal{V}$-face algebra $\mathfrak{H}$ is called *compact* if $\mathfrak{H}$ is spanned by entries of unitary matrix corepresentations (cf. [29, 17]). For each costar $\mathcal{V}$-face algebra $\mathfrak{H}$, its dual $\mathfrak{H}^*$ becomes a $*$-algebra via

$$\langle X^*, a\rangle = \overline{\langle X, a^\times\rangle} \quad (X \in \mathfrak{H}^*, \mathfrak{a} \in \mathfrak{H}). \tag{4.4}$$

When $\mathfrak{H}$ is a Hopf $\mathcal{V}$-face algebra, we also set $a^* = S(a^\times)$ and $X^\times = S^*(X)^*$ for each $a \in \mathfrak{H}$ and $X \in \mathfrak{H}^*$. These operations satisfy

$$(a^*)^* = a, \quad (X^\times)^\times = X \quad (a \in \mathfrak{H}, \mathfrak{X} \in \mathfrak{H}^*). \tag{4.5}$$

For each compact Hopf $\mathcal{V}$-face algebra $\mathfrak{H}$, its *Woronowicz functional* [29, 17] $\mathcal{Q} \in \mathfrak{H}^*$ is a unique group-like functional such that

$$\mathcal{Q}\mathcal{X}\mathcal{Q}^{-\infty} = (\mathcal{S}^{\in})^*(\mathcal{X}) \quad (\mathcal{X} \in \mathfrak{H}^*), \tag{4.6}$$

$$\sum_p \mathcal{Q}(\sqcup_{\surd\surd}) = \sum_{\surd} \mathcal{Q}^{-\infty}(\sqcup_{\surd\surd}), \tag{4.7}$$

and that $[\mathcal{Q}(\sqcup_{\surd}{}_{\mathrm{II}})]$ is a positive matrix, where $[t_{pq}]$ denotes an arbitrary unitary matrix corepresentation. The functional $\mathcal{Q}$ also satisfies

$$\mathcal{Q}^* = \mathcal{Q}, \quad \mathcal{Q}^\times = \mathcal{Q}^{-\infty}. \tag{4.8}$$

Let $(\mathfrak{H}, \times)$ be a costar $\mathcal{V}$-face algebra and M a finite-dimensional $\mathfrak{H}$-comodule equipped with a Hilbert space structure $(\,|\,)$. We say that $(M, (\,|\,))$ is *unitary* if

$$\sum_{(u)} (u_{(0)} \,|\, v)\, u_{(1)} = \sum_{(v)} (u \,|\, v_{(0)})\, v_{(1)}^\times \quad (u, v \in M). \tag{4.9}$$

We note that $(M, (\,|\,))$ is unitary if and only if (3.11) gives a $*$-representation of $\mathfrak{H}^*$. A costar $\mathcal{V}$-face algebra $(\mathfrak{H}, \times)$ is compact if and only if every finite-dimensional $\mathfrak{H}$-comodule is unitary for some $(\,|\,)$.

Proposition 4.1. *Let $\mathfrak{H}$ be a compact $\mathcal{V}$-face algebra.* (1) *The unit comodule $\mathbb{C}\mathcal{V}$ becomes a unitary comodule via $(\lambda \,|\, \mu) = \delta_{\lambda\mu}$.* (2) *For each unitary comodules M and N, $M \bar{\otimes} N$ becomes a unitary comodule via $(u \otimes v \,|\, u' \otimes v') = (u \,|\, u')(v \,|\, v')$ $(u \in M(\lambda, \nu), v \in N(\nu, \mu), u' \in M(\lambda', \nu'), v' \in N(\nu', \mu'))$.* (3) *If $\mathfrak{H}$ has an antipode, then the left dual $M^\vee$ becomes a unitary comodule via*

$$(u \,|\, v) = (\mathcal{Q}\Upsilon^{-\infty}(\sqsubseteq) \,|\, \Upsilon^{-\infty}(\sqcap)), \tag{4.10}$$

where $\Upsilon \colon M \to M^\vee$ denotes the antilinear isomorphism defined by

$$\langle \Upsilon(u), v \rangle = (v \,|\, u) \quad (u, v \in M). \tag{4.11}$$

Proof. The proof of Part (1) and Part (2) is straightforward. Using the second equality of (4.5), we obtain $(S^2)^*(X^\times) = S^*(X^*)$. Using this together with (4.6) and

$$X\Upsilon(u) = \Upsilon(X^\times u) \quad (u \in M, X \in \mathfrak{H}^*), \tag{4.12}$$

we obtain

$$\begin{aligned} (u \,|\, Xv) &= (S^*(X^*)\mathcal{Q}\Upsilon^{-\infty}(\sqsubseteq) \,|\, \Upsilon^{-\infty}(\sqcap)) \\ &= (\mathcal{Q}\Upsilon^{-\infty}(\sqsubseteq) \,|\, (\mathcal{X}^*)^\times \Upsilon^{-\infty}(\sqcap)) \\ &= (X^* u \,|\, v) \end{aligned}$$

for each $u, v \in M^\vee$, as required. □

By the proposition above, the category $\mathbf{Com}_{\mathfrak{H}}^{\mathbf{fu}}$ of all finite-dimensional unitary comodules of a compact (Hopf) $\mathcal{V}$-face algebra $\mathfrak{H}$ becomes a (rigid) monoidal category. Let M and N be unitary $\mathfrak{H}$-comodules and $f \colon M \to N$ an $\mathfrak{H}$-comodule map. We define a linear map $\bar{f} \colon N \to M$ by $(\overline{f}(n) \,|\, m) = (n \,|\, f(m))$. We have

$$\overline{\overline{f}} = f, \quad \overline{f \circ g} = \overline{g} \circ \overline{f}, \quad \overline{f \bar{\otimes} g} = \overline{f} \bar{\otimes} \overline{g}, \tag{4.13}$$

$$\overline{f + g} = \overline{f} + \overline{g}, \quad \overline{cf} = \overline{c}\overline{f} \quad (c \in \mathbb{C}). \tag{4.14}$$

Proposition 4.2. *For each compact Hopf $\mathcal{V}$-face algebra $\mathfrak{H}$ and its comodule M, we have*

$$\overline{d_M} = ((I_{M^\wedge} \circ \mathcal{Q}) \bar{\otimes} \mathrm{id_M}) \circ b_{M^\wedge}, \tag{4.15}$$

$$\overline{b_M} = d_{M^\wedge} \circ (\mathrm{id_M} \bar{\otimes} ((\mathrm{I_{M^\wedge}} \circ \mathcal{Q})^{-\infty}). \tag{4.16}$$

Proof. Using (4.8) and (4.12), we obtain

$$(\mathbf{q}^\vee \,|\, \mathcal{Q}\mathbf{p}^\vee) = \overline{\langle \mathbf{p}^\vee,\, \Upsilon^{-1}(\mathbf{q}^\vee)\rangle}, \quad (\Upsilon(\mathbf{q}) \,|\, \mathbf{r}^\vee) = \overline{\langle \mathbf{r}^\vee,\, \mathcal{Q}\mathbf{q}\rangle}, \tag{4.17}$$

where $\{\mathbf{p}\}$ and $\{\mathbf{p}^\vee\}$ is as in (3.8). Using the first equality of (4.17), we obtain

$$(\mathbf{q}^\vee \otimes \mathbf{r} \,|\, ((\mathbf{I}_{\mathbf{M}^\wedge} \circ \mathcal{Q})\bar{\otimes}\mathrm{id}_{\mathrm{M}}) \circ \mathrm{b}_{\mathrm{M}^\wedge}(\lambda)) = \sum_{\mu \in \mathcal{V}} \sum_{\mathbf{p} \in \mathcal{M}_{\mu\lambda}} (\mathbf{q}^\vee \otimes \mathbf{r} \,|\, \mathcal{Q}\mathbf{p}^\vee \otimes \mathbf{p})$$

$$= \sum_{\mu \in \mathcal{V}} \sum_{\mathbf{p} \in \mathcal{M}_{\mu\lambda}} \big(\mathbf{r} \,|\, \langle \mathbf{p}^\vee, \Upsilon^{-1}(\mathbf{q}^\vee)\rangle \mathbf{p}\big) = \delta_{\lambda \mathfrak{r}(\mathbf{q})}\, \delta_{\mathbf{qr}},$$

where the first equality follows from (3.10). This proves (4.15). Similarly, (4.16) follows from the second equality of (4.17). □

Let $\mathfrak{H}$ be a costar CQT $\mathcal{V}$-face algebra. We say that $\mathfrak{H}$ is of *unitary type* if

$$\mathcal{R}^+(a^\times, b^\times) = \overline{\mathcal{R}^-(a,b)} \quad (a, b \in \mathfrak{H}). \tag{4.18}$$

In this case, the Drinfeld functionals of $\mathfrak{H}$ satisfy

$$\mathcal{U}_\infty^* = \mathcal{U}_\in, \quad \mathcal{U}_\in^* = \mathcal{U}_\infty. \tag{4.19}$$

Remark 4.1. When $q > 0$, the function algebras of the usual quantum groups (such as $\mathrm{Fun}(\mathrm{SL_q(N)})$, $\mathrm{Fun}(\mathrm{Sp_q(2N)})$) are both CQT and compact. However, they are not of unitary type but rather of "Hermitian type."

Proposition 4.3 ([22]). *For each compact CQT Hopf $\mathcal{V}$-face algebra $\mathfrak{H}$ of unitary type, its Woronowicz functional is a modified ribbon functional of $\mathfrak{H}$. Moreover, the corresponding ribbon functional $\mathcal{V}_\mathcal{Q}$ satisfies*

$$\mathcal{V}_\mathcal{Q}^* = \mathcal{V}_\mathcal{Q}^{-\infty}. \tag{4.20}$$

We call $\mathcal{V}_\mathcal{Q}$ the *canonical ribbon functional* of $(\mathfrak{H}, \times)$. We note that the expression $\mathcal{U}_\infty = \mathcal{V}_\mathcal{Q}\mathcal{Q}$ gives the "polar decomposition" of $\mathcal{U}_\infty$.

Let $\mathfrak{H}$ be a compact CQT Hopf $\mathcal{V}$-face algebra of unitary type, equipped with the canonical ribbon functional. By (4.18) and (4.20), we have

$$\overline{c_{MN}} = (c_{MN})^{-1}, \quad \overline{\theta_M} = \theta_M^{-1}. \tag{4.21}$$

Moreover, by (3.15), (3.16), (4.16), we have

$$\overline{b_M} = d_M \circ c_{MM^\vee} \circ (\theta_M \bar{\otimes} \mathrm{id}_{\mathrm{M}^\vee}). \tag{4.22}$$

Furthermore, by (3.14) and the defining properties of $\mathcal{Q}$, we have

$$\mathrm{Tr_q}(\mathrm{f}\bar{\mathrm{f}}) = \frac{1}{\mathrm{card}(\mathcal{V})} \mathrm{Tr}(\bar{\mathrm{f}}\mathcal{Q}\{) > \prime \tag{4.23}$$

for every $f \neq 0$. Thus, the category $\mathbf{Com}_{\mathfrak{H}}^{\mathbf{fu}}$ is a unitary ribbon category (cf. [40]).

Example 4.1. Let $L = 1$ and $\mathfrak{S}(A_{N-1}; t)_\epsilon$ as in Example 2.1. Then $\mathfrak{S}(A_{N-1}; t)_\epsilon$ is a compact CQT Hopf $\mathcal{V}$-face algebra of unitary type with costar structure $e^i_j(m)^\times = e^j_i(m)$. Its Woronowicz functional agrees with the counit ε.

5. FLAT FACE MODELS

Let $\mathcal{G}$ be a finite oriented graph with set of vertices $\mathcal{V}$. We say that a quadruple $\left(\mathbf{r}\,{}^{\mathbf{p}}_{\mathbf{q}}\,\mathbf{s}\right)$ or a diagram

$$\begin{tikzcd} \lambda \arrow[r, "\mathbf{p}"] \arrow[d, "\mathbf{r}"'] & \mu \arrow[d, "\mathbf{s}"] \\ \nu \arrow[r, "\mathbf{q}"] & \xi \end{tikzcd} \tag{5.1}$$

is a *face* if $\mathbf{p}, \mathbf{q}, \mathbf{r}, \mathbf{s} \in \mathcal{G}^{\mathbf{1}}$ and

$$\mathfrak{s}(\mathbf{p}) = \lambda = \mathfrak{s}(\mathbf{r}), \quad \mathfrak{r}(\mathbf{p}) = \mu = \mathfrak{s}(\mathbf{s}), \quad \mathfrak{r}(\mathbf{r}) = \nu = \mathfrak{s}(\mathbf{q}), \quad \mathfrak{r}(\mathbf{q}) = \xi = \mathfrak{r}(\mathbf{s}). \tag{5.2}$$

When $\mathcal{G}$ has no multiple edge, we also write $\left(\mathbf{r}\,{}^{\mathbf{p}}_{\mathbf{q}}\,\mathbf{s}\right) = \left(\begin{smallmatrix}\lambda & \mu \\ \nu & \xi\end{smallmatrix}\right)$. We say that $(\mathcal{G},)$ is a ($\mathcal{V}$-)*face model* over a field $\mathbb{K}$ if w is a map which assigns a number $w\!\left[\mathbf{r}\,{}^{\mathbf{p}}_{\mathbf{q}}\,\mathbf{s}\right] \in \mathbb{K}$ to each face $\left(\mathbf{r}\,{}^{\mathbf{p}}_{\mathbf{q}}\,\mathbf{s}\right)$ of $\mathcal{G}$. We call w *Boltzmann weight* of $(\mathcal{G},)$. For convenience, we set $w\!\left[\mathbf{r}\,{}^{\mathbf{p}}_{\mathbf{q}}\,\mathbf{s}\right] = 0$ unless $\left(\mathbf{r}\,{}^{\mathbf{p}}_{\mathbf{q}}\,\mathbf{s}\right)$ is a face. For a face model $(\mathcal{G},)$, we identify w with the linear operator on $\mathbb{K}\mathcal{G} = \bigoplus_{\mathbf{p}\in\mathcal{G}} \mathbb{K}\mathbf{p}$ given by

$$w(\mathbf{p}\cdot\mathbf{q}) = \sum_{\mathbf{r}\cdot\mathbf{s}\in\mathcal{G}^{\mathbf{2}}} \mathbf{w}\!\left[\mathbf{r}\,{}^{\mathbf{p}}_{\mathbf{s}}\,\mathbf{q}\right] \mathbf{r}\cdot\mathbf{s} \quad (\mathbf{p}\cdot\mathbf{q} \in \mathcal{G}). \tag{5.3}$$

For $m \geq 2$ and $1 \leq i < m$, we define an operator $w_i = w_{i/m}$ on $\mathbb{K}\mathcal{G}$ by $w_{i/m}(\mathbf{p}\cdot\mathbf{q}\cdot\mathbf{r}) = \mathbf{p} \otimes \mathbf{w}(\mathbf{q}) \otimes \mathbf{r}$ ($\mathbf{p} \in \mathcal{G}^{-}, \mathbf{q} \in \mathcal{G}, \mathbf{r} \in \mathcal{G}^{--}$), where we identify $(\mathbf{p_1}, \ldots, \mathbf{p_m}) \in \mathcal{G}$ with $\mathbf{p_1} \otimes \ldots \otimes \mathbf{p_m} \in (\mathbb{K}\mathcal{G})^{\otimes}$.

A face model is called *invertible* if w is invertible as an operator on $\mathbb{K}\mathcal{G}$. An invertible face model is called *star-triangular* (or *Yang-Baxter*) if w satisfies the braid relation $w_1 w_2 w_1 = w_2 w_1 w_2$ in $\mathrm{End}(\mathbb{K}\mathcal{G})$.

For a star-triangular face model $(\mathcal{G},)$, the operators $w_{i/m}$ $(1 \leq i < m)$ define an action of the m-string braid group $\mathfrak{B}_{\mathfrak{m}}$ on $\mathbb{K}\mathcal{G}_{\lambda\mu}$ for each $m \geq 2$ and $\lambda, \mu \in \mathcal{V}$. The following proposition gives a face version of the FRT construction.

Proposition 5.1 ([35, 30, 13, 19])**.** *Let* $(\mathcal{G},)$ *be a* $\mathcal{V}$*-face model and* $\mathfrak{H}(\mathcal{G})$ *as in* § *1. Let* $\mathfrak{I}$ *be an ideal of* $\mathfrak{H}(\mathcal{G})$ *generated by the following elements*:

$$\sum_{\mathbf{r}\cdot\mathbf{s}\in\mathcal{G}^{\mathbf{2}}} w\!\left[\mathbf{r}\,{}^{\mathbf{p}}_{\mathbf{s}}\,\mathbf{q}\right] e\begin{pmatrix}\mathbf{a}\cdot\mathbf{b}\\ \mathbf{r}\cdot\mathbf{s}\end{pmatrix} - \sum_{\mathbf{c}\cdot\mathbf{d}\in\mathcal{G}^{\mathbf{2}}} w\!\left[\mathbf{a}\,{}^{\mathbf{c}}_{\mathbf{b}}\,\mathbf{d}\right] e\begin{pmatrix}\mathbf{c}\cdot\mathbf{d}\\ \mathbf{p}\cdot\mathbf{q}\end{pmatrix} \quad (\mathbf{p}\cdot\mathbf{q},\ \mathbf{a}\cdot\mathbf{b} \in \mathcal{G}). \tag{5.4}$$

Then $\mathfrak{I}$ *is a coideal of* $\mathfrak{H}(\mathcal{G})$ *and the quotient* $\mathfrak{A}(w) := \mathfrak{H}(\mathcal{G})/\mathfrak{I}$ *becomes a* $\mathcal{V}$*-face algebra. If* $(\mathcal{G},)$ *is star-triangular, then there exist unique bilinear pairings* $\mathcal{R}^{\pm}$ *on* $\mathfrak{A}(w)$ *such that* $(\mathfrak{A}(w), \mathcal{R}^{\pm})$ *is a CQT* $\mathcal{V}$*-face algebra and that*

$$\mathcal{R}^{+}\left(e\begin{pmatrix}\mathbf{p}\\ \mathbf{q}\end{pmatrix}, e\begin{pmatrix}\mathbf{r}\\ \mathbf{s}\end{pmatrix}\right) = w\!\left[\mathbf{r}\,{}^{\mathbf{q}}_{\mathbf{p}}\,\mathbf{s}\right] \tag{5.5}$$

for each $\mathbf{p}, \mathbf{q}, \mathbf{r}, \mathbf{s} \in \mathcal{G}^{\mathbf{1}}$.

We denote the image of $e\binom{\mathbf{p}}{\mathbf{q}}$ by the projection $\mathfrak{H}(\mathcal{G}) \to \mathfrak{A}()$ again by $e\binom{\mathbf{p}}{\mathbf{q}}$. Then $\mathfrak{A}^{\mathfrak{m}}(\mathfrak{w}) := \sum_{\mathbf{p},\mathbf{q}\in\mathcal{G}} \mathbb{K}\binom{\mathbf{p}}{\mathbf{q}}$ becomes a subcoalgebra of $\mathfrak{A}(w)$ for each $m \geq 0$. As the usual FRT construction (cf. [12, Proposition 2.1]), we have the following.

Proposition 5.2. *For each star-triangular face model* $(\mathcal{G},)$*, we have* $\mathfrak{A}^{\mathfrak{m}}(\mathfrak{w})^* \cong \mathrm{Hom}_{\mathfrak{B}_{\mathfrak{m}}}(\mathbb{K}\mathcal{G})$ $(m \geq 2)$ *as* $\mathbb{K}$*-algebras.*

We say that $\left(\mathbf{r}\,{}^{\mathbf{p}}_{\mathbf{q}}\,\mathbf{s}\right)$ or (5.1) is a *boundary condition* of size $m \times n$ if $\mathbf{p}, \mathbf{q} \in \mathcal{G}^{\mathbf{n}}, \mathbf{r}, \mathbf{s} \in \mathcal{G}^{\mathbf{m}}$ and the relation (5.2) is satisfied for some λ, μ, ν, ξ. For a face model $(\mathcal{G},)$, we define its *partition function* to be an extension $w : \{$ boundary conditions of size $m \times n;\ m, n \geq 1\} \to \mathbb{K}$ of the map w which is determined by the following two recursion relations:

$$w\begin{bmatrix} & \mathbf{p}\cdot\mathbf{p}' & \\ \mathbf{r} & & \mathbf{s} \\ & \mathbf{q}\cdot\mathbf{q}' & \end{bmatrix} = \sum_{\mathbf{a}\in\mathcal{G}^{\mathbf{m}}} w\begin{bmatrix} & \mathbf{p} & \\ \mathbf{r} & & \mathbf{a} \\ & \mathbf{q} & \end{bmatrix} w\begin{bmatrix} & \mathbf{p}' & \\ \mathbf{a} & & \mathbf{s} \\ & \mathbf{q}' & \end{bmatrix}, \tag{5.6}$$

$$w\begin{bmatrix} & \mathbf{p} & \\ \mathbf{r}\cdot\mathbf{r}' & & \mathbf{s}\cdot\mathbf{s}' \\ & \mathbf{q} & \end{bmatrix} = \sum_{\mathbf{a}\in\mathcal{G}^{\mathbf{n}}} w\begin{bmatrix} & \mathbf{p} & \\ \mathbf{r} & & \mathbf{s} \\ & \mathbf{a} & \end{bmatrix} w\begin{bmatrix} & \mathbf{a} & \\ \mathbf{r}' & & \mathbf{s}' \\ & \mathbf{q} & \end{bmatrix} \tag{5.7}$$

$$\left(\mathbf{p},\mathbf{q}\in\mathcal{G}^{\mathbf{n}},\mathbf{p}',\mathbf{q}'\in\mathcal{G}^{\mathbf{n}'},\mathbf{r},\mathbf{s}\in\mathcal{G}^{\mathbf{m}},\mathbf{r}',\mathbf{s}'\in\mathcal{G}^{\mathbf{m}'}\right).$$

Also, we set $w\begin{bmatrix} & \mathbf{p} & \\ \mathbf{r} & & \mathbf{s} \\ & \mathbf{q} & \end{bmatrix} = \delta_{\mathbf{pq}}$ (respectively $w\begin{bmatrix} & \mathbf{p} & \\ \mathbf{r} & & \mathbf{s} \\ & \mathbf{q} & \end{bmatrix} = \delta_{\mathbf{rs}}$) if $\mathbf{r},\mathbf{s}\in\mathcal{G}^{\mathbf{0}}$ (respectively $\mathbf{p},\mathbf{q}\in\mathcal{G}^{\mathbf{0}}$). With this notation, the relation (5.5) holds for every star-triangular face model $(\mathcal{G},)$ and $\mathbf{p},\mathbf{q}\in\mathcal{G}^{\mathbf{n}},\mathbf{r},\mathbf{s}\in\mathcal{G}^{\mathbf{m}}(\mathbf{m},\mathbf{n}\geq\mathbf{0})$.

Next, we recall the notion of flat face model [18], which is a variant of A. Ocneanu's notion of flat biunitary connection (cf. [33]). Let $(\mathcal{G},)$ be an invertible face model with a fixed vertex $*\in\mathcal{V}=\mathcal{G}$. We assume that $\boldsymbol{\Lambda}^m_{\mathcal{G}}\neq\emptyset$ for each $m\geq 0$ and that $\mathcal{V}=\bigcup_{\geq}\mathcal{V}()$, where $\boldsymbol{\Lambda}^m_{\mathcal{G}}=\boldsymbol{\Lambda}^m_{\mathcal{G}*}$ and $\mathcal{V}()=\mathcal{V}()_{\mathcal{G}*}$ are defined by

$$\boldsymbol{\Lambda}_{\mathcal{G}} = \left\{(\lambda,m)\in\mathcal{V}\times\mathbb{Z}_{\geq\nvdash}\,\middle|\,\mathcal{G}_{*\lambda}\neq\emptyset\right\}, \tag{5.8}$$

$$\boldsymbol{\Lambda}^m_{\mathcal{G}} = \boldsymbol{\Lambda}_{\mathcal{G}}\cap(\mathcal{V}\times\{\}), \tag{5.9}$$

$$\mathcal{V}() = \left\{\lambda\in\mathcal{V}\,\middle|\,(\lambda,)\in\boldsymbol{\Lambda}_{\mathcal{G}}\right\}. \tag{5.10}$$

For each $m\geq 0$, we define the algebra $\mathrm{Str^m}(\mathcal{G},*)$ by

$$\mathrm{Str^m}(\mathcal{G},*) = \prod_{\lambda\in\mathcal{V}()}\mathrm{End}(\mathbb{K}\mathcal{G}_{*\lambda}) \tag{5.11}$$

and call it *string algebra* of $(\mathcal{G}, ,*)$. For each $m,n\geq 0$, we define the algebra map $\iota_{mn}\colon\mathrm{Str^m}(\mathcal{G},*)\to\mathrm{Str^{m+n}}(\mathcal{G},*)$ by $\iota_{mn}(x)(\mathbf{p}\cdot\mathbf{q})=\mathbf{xp}\otimes\mathbf{q}$ $(\mathbf{p}\in\mathcal{G}_{*\lambda},\mathbf{q}\in\mathcal{G}^{\backslash}_{\lambda\mu})$. For each $1\leq i<m$, we define the element $\overset{*}{w}_i=\overset{*}{w}_{i/m}$ of $\mathrm{Str^m}(\mathcal{G},*)$ to be the restriction of $w_{i/m}$ on $\mathbb{K}\mathcal{G}_{*-}$, where $\mathcal{G}_{*-}=\coprod_{\lambda\in\mathcal{V}}\mathcal{G}_{*\lambda}$. We say that $(\mathcal{G}, ,*)$ is a *flat face model* if

$$\iota_{mn}(x)\overset{*}{w}_{nm}\iota_{nm}(y)\overset{*}{w}{}^{-1}_{nm} = \overset{*}{w}_{nm}\iota_{nm}(y)\overset{*}{w}{}^{-1}_{nm}\iota_{mn}(x) \tag{5.12}$$

for each $x\in\mathrm{Str^m}(\mathcal{G},*)$ and $y\in\mathrm{Str^n}(\mathcal{G},*)$ $(m,n\geq 0)$, where $\overset{*}{w}_{mn}\in\mathrm{Str^{m+n}}(\mathcal{G},*)$ is defined by

$$\overset{*}{w}_{mn} = (\overset{*}{w}_n\overset{*}{w}_{n+1}\cdots\overset{*}{w}_{m+n-1})(\overset{*}{w}_{n-1}\overset{*}{w}_n\cdots\overset{*}{w}_{m+n-2})\cdots(\overset{*}{w}_1\overset{*}{w}_2\cdots\overset{*}{w}_m). \tag{5.13}$$

For each flat $\mathcal{V}$-face model $(\mathcal{G}, ,*)$, $n\geq 0$ and $\lambda,\mu\in\mathcal{V}$, there exists a unique left action Γ of $\mathrm{Str^n}(\mathcal{G},*)$ on $\mathbb{K}\mathcal{G}^{\backslash}_{\lambda\mu}$ such that

$$\mathbf{p}\otimes(\Gamma(\mathbf{x})\mathbf{q}) = \overset{*}{\mathbf{w}}_{\mathbf{nm}}\iota_{\mathbf{nm}}(\mathbf{x})\overset{*}{\mathbf{w}}{}^{\mathbf{-1}}_{\mathbf{nm}}(\mathbf{p}\cdot\mathbf{q}) \tag{5.14}$$

for each $m\geq 0$, $\mathbf{p}\in\mathcal{G}_{*\lambda},\mathbf{q}\in\mathcal{G}^{\backslash}_{\lambda\mu}$ and $x\in\mathrm{Str^n}(\mathcal{G},*)$. Using this action, we define *costring algebra*

$$\mathrm{Cost(w,*)} = \bigoplus_{\mathrm{m}\geq 0}\mathrm{Cost^m(w,*)} \tag{5.15}$$

to be the quotient $\mathcal{V}$-face algebra of $\bigoplus_{m\geq 0}\mathrm{End}_{\mathbb{K}}(\mathbb{K}\mathcal{G})^*\cong\mathfrak{H}(\mathcal{G})$ given by

$$\mathrm{Cost^m(w,*)} = \mathrm{End}_{\mathrm{Str^m}(\mathcal{G},*)}(\mathbb{K}\mathcal{G})^*. \tag{5.16}$$

For each $\lambda,\mu\in\mathcal{V}$ and $(\nu,m)\in\boldsymbol{\Lambda}_{\mathcal{G}}$, we define the non-negative integer $N^{\mu}_{\lambda\nu}(m)$ by the irreducible decomposition of $\mathbb{K}\mathcal{G}_{\lambda\mu}$:

$$[\mathbb{K}\mathcal{G}_{\lambda\mu}] = \sum_{\nu\in\mathcal{V}()}\mathcal{N}^{\mu}_{\lambda\nu}()[\mathbb{K}\mathcal{G}_{*\nu}], \tag{5.17}$$

where, for each $\mathrm{Str}^{\mathrm{m}}(\mathcal{G}, *)$-module V, $[V]$ denotes the element of the Grothendieck group $K_0(\mathrm{Str}^{\mathrm{m}}(\mathcal{G}, *))$ corresponding to V (see e.g. [4, §5.1]). We call $N^{\mu}_{\lambda\nu}(m)$ *fusion rules* of $(\mathcal{G}, , *)$,

Theorem 5.3 ([18])**.** *Let* $(\mathcal{G}, , *)$ *be a flat* $\mathcal{V}$*-face model with fusion rule* $N^{\mu}_{\lambda\nu}(m)$. (1) *For each* $(\lambda, m) \in \mathbf{\Lambda}_{\mathcal{G}}$*, up to isomorphism there exists a unique right* $\mathrm{Cost}^{\mathrm{m}}(\mathrm{w}, *)$*-comodule* $L_{(\lambda,m)}$ *such that*

$$\dim L_{(\lambda,m)}(*, \mu) = \delta_{\lambda\mu} \tag{5.18}$$

for each $\mu \in \mathcal{V}$*. As coalgebras, we have*

$$\mathrm{Cost}^{\mathrm{m}}(\mathrm{w}, *) \cong \bigoplus_{\lambda \in \mathcal{V}()} \mathrm{End}(\mathrm{L}_{(\lambda,\mathrm{m})})^*. \tag{5.19}$$

(2) *In the corepresentation ring* $K_0(\mathbf{Com}^{\mathbf{f}}_{\mathrm{Cost}(\mathrm{w},*)})$*, we have*

$$[L_{(*,0)}] = 1, \tag{5.20}$$

$$[L_{(\lambda,m)}][L_{(\mu,n)}] = \sum_{\nu \in \mathcal{V}(+\backslash)} N^{\nu}_{\lambda\mu}(n)[L_{(\nu,m+n)}]. \tag{5.21}$$

Moreover, for each $\mathrm{Cost}^{\mathrm{m}}(\mathrm{w}, *)$*-comodule* M*, we have*

$$[M] = \sum_{\lambda \in \mathcal{V}()} \dim\left(M(*, \lambda)\right)[L_{(\lambda,m)}]. \tag{5.22}$$

(3) *We have*

$$\dim\left(L_{(\nu,m)}(\lambda, \mu)\right) = N^{\mu}_{\lambda\nu}(m). \tag{5.23}$$

Lemma 5.4. *For each flat star-triangular face model, we have*

$$\Gamma(\overset{*}{w}_{i/n}) = w_{i/n} \quad (n \geq 2,\ 1 \leq i < n). \tag{5.24}$$

Proof. By the braid relation, we have

$$\overset{*}{w}_{nm}\overset{*}{w}_{i/n+m}\overset{*}{w}_{nm}^{-1} = \overset{*}{w}_{i+m/n+m} \tag{5.25}$$

for each $1 \leq i < n$. Using this together with $\iota_{nm}(\overset{*}{w}_{i/n}) = \overset{*}{w}_{i/n+m}$, we obtain

$$\mathbf{p} \otimes (\Gamma(\overset{*}{\mathbf{w}}_{\mathbf{i/n}})\mathbf{q}) = \mathbf{p} \otimes \mathbf{w}_{\mathbf{i/n}}\mathbf{q} \tag{5.26}$$

for each $m \geq 0$, $\mathbf{p} \in \mathcal{G}_{*\lambda}$ and $\mathbf{q} \in \mathcal{G}^{\backslash}_{\lambda\mu}$ as required. □

Proposition 5.5. *Let* $(\mathcal{G},)$ *be a star-triangular* $\mathcal{V}$*-face model with a fixed vertex* $* \in \mathcal{V}$*. Then* $(\mathcal{G}, , *)$ *is flat if* $\mathbb{K}\mathcal{G}_{*\lambda}$ *is an absolutely irreducible* $\mathfrak{B}_{\mathfrak{m}}$*-module for each* $(\lambda, m) \in \mathbf{\Lambda}_{\mathcal{G}}$*. In this case, we have* $\mathrm{Cost}(\mathrm{w}, *) = \mathfrak{A}(w)$ *as quotients of* $\mathfrak{H}(\mathcal{G})$*.*

Proof. Using (5.25), we see that (5.12) holds for every $x \in \mathrm{Str}^{\mathrm{m}}(\mathcal{G}, *)$ and $y = \overset{*}{w}_i$ $(1 \leq i < n)$. Hence $(\mathcal{G}, , *)$ is flat if $\mathrm{Str}^{\mathrm{m}}(\mathcal{G}, *) = \langle {}^*, \ldots, {}^*{}_- \rangle$ for each $m > 1$. The second assertion follows from Proposition 5.2 and the lemma above. □

6. $SU(N)_L$-SOS MODELS

In order to construct the algebras $\mathfrak{S}(A_{N-1}; t)_\epsilon$, we first recall $SU(N)_L$-SOS models (without spectral parameter) [23], which are equivalent to H. Wenzl's representations of Iwahori-Hecke algebras (cf. [45]) and also, the monodromy representations of the braid group arising from conformal field theory (cf. A. Tsuchiya and Y. Kanie [39]). Let $N \geq 2$ and $L \geq 2$ be integers. For each $1 \leq i \leq N$, we define the vector $\hat{i} \in \mathbb{R}^{\mathbb{N}}$ by $\hat{1} = (1 - 1/N, -1/N, \ldots, -1/N)$, …, $\hat{N} = (-1/N, \ldots, -1/N, 1 - 1/N)$. Let $\mathcal{V} = \mathcal{V}_{\mathcal{NL}}$ be the subset of $\mathbb{R}^{\mathbb{N}}$ given by

$$\mathcal{V}_{NL} = \{\lambda_1\hat{1} + \cdots \lambda_N\hat{N} \mid \lambda_1, \ldots, \lambda_N \in \mathbb{Z}, \mathbb{L} \geq \lambda_{\nVdash} \geq \cdots \geq \lambda_{\mathbb{N}} = \nvdash\}. \tag{6.1}$$

For $\lambda \in \mathcal{V}$, we define integers $\lambda_1, \ldots, \lambda_N$ and $|\lambda|$ by $\lambda = \sum_i \lambda_i\hat{i}$, $\lambda_N = 0$ and $|\lambda| = \sum_i \lambda_i$. For $m \geq 0$, we define a subset $\mathcal{G}^m$ of $\mathcal{V}^{m+1}$ by

$$\mathcal{G}^m = \mathcal{V}^{m+1} \cap \{\mathbf{p} = (\lambda \,|\, \mathbf{i_1}, \ldots, \mathbf{i_m}) \mid \lambda \in \mathcal{V},\ \leq, \ldots,\ \leq \mathcal{N}\}, \tag{6.2}$$

where for $\lambda \in \mathbb{R}^{\mathbb{N}}$ and $1 \leq i_1, \ldots, i_m \leq N$, we set

$$(\lambda \,|\, i_1, \ldots, i_m) = (\lambda, \lambda + \hat{i}_1, \ldots, \lambda + \hat{i}_1 + \cdots + \hat{i}_m). \tag{6.3}$$

Then $(\mathcal{V}, \mathcal{G})$ defines an oriented graph $\mathcal{G} = \mathcal{G}_{\mathcal{N},\mathcal{L}}$ and $\mathcal{G}^m$ is identified with the set of paths of $\mathcal{G}$ of length m. For $\mathbf{p} = (\lambda \,|\, \mathbf{i}, \mathbf{j})$, we set $\mathbf{p}^\dagger = (\lambda \,|\, j, i)$ and

$$d(\mathbf{p}) = \lambda_{\mathbf{i}} - \lambda_{\mathbf{j}} + \mathbf{j} - \mathbf{i}. \tag{6.4}$$

We define subsets $\mathcal{G}^2[\rightarrow]$, $\mathcal{G}^2[\downarrow]$ and $\mathcal{G}^2[\searrow]$ of $\mathcal{G}^2$ by

$$\mathcal{G}^2[\rightarrow] = \{\mathbf{p} \in \mathcal{G}^{\mathbf{2}} \mid \mathbf{p}^\dagger = \mathbf{p}\}, \tag{6.5}$$

$$\mathcal{G}^2[\downarrow] = \{\mathbf{p} \in \mathcal{G}^{\mathbf{2}} \mid \mathbf{p}^\dagger \notin \mathcal{G}\}, \tag{6.6}$$

$$\mathcal{G}^2[\searrow] = \{\mathbf{p} \in \mathcal{G}^{\mathbf{2}} \mid \mathbf{p} \neq \mathbf{p}^\dagger \in \mathcal{G}\}. \tag{6.7}$$

Let $t \in \mathbb{C}$ be a primitive $2(N+L)$-th root of 1. Let ϵ be either 1 or -1 and ζ a nonzero parameter. We define a face model $(\mathcal{G}, {}_{\mathcal{N},,\epsilon}) = (\mathcal{G}_{\mathcal{N},\mathcal{L}}, {}_{\mathcal{N},,\epsilon,\zeta})$ by setting

$$w_{N,t,\epsilon}\begin{bmatrix} \lambda & \lambda+\hat{i} \\ \lambda+\hat{i} & \lambda+\hat{i}+\hat{j} \end{bmatrix} = -\zeta^{-1}t^{-d(\mathbf{p})}\frac{1}{[d(\mathbf{p})]}, \tag{6.8}$$

$$w_{N,t,\epsilon}\begin{bmatrix} \lambda & \lambda+\hat{i} \\ \lambda+\hat{j} & \lambda+\hat{i}+\hat{j} \end{bmatrix} = \zeta^{-1}\epsilon\frac{[d(\mathbf{p})-\mathbf{1}]}{[\mathbf{d}(\mathbf{p})]}, \tag{6.9}$$

$$w_{N,t,\epsilon}\begin{bmatrix} \lambda & \lambda+\hat{k} \\ \lambda+\hat{k} & \lambda+2\hat{k} \end{bmatrix} = \zeta^{-1}t \tag{6.10}$$

for each $\mathbf{p} = (\lambda \,|\, \mathbf{i}, \mathbf{j}) \in \mathcal{G}[\searrow] \amalg \mathcal{G}[\downarrow]$ and $(\lambda \,|\, k, k) \in \mathcal{G}[\rightarrow]$, where $[n] = (t^n - t^{-n})/(t - t^{-1})$ for each $n \in \mathbb{Z}$. We call $(\mathcal{G}, {}_{\mathcal{N},,\epsilon})$ *$SU(N)_L$-SOS model* (without spectral parameter) [23]. It is known that $(\mathcal{G}, {}_{\mathcal{N},,\epsilon})$ is star-triangular. Moreover, H. Wenzl [45] showed that $\mathbb{C}\mathcal{G}_\lambda$ is an irreducible $\mathfrak{B}_{\mathfrak{m}}$-module for each $m \geq 0$ and $\lambda \in \Lambda^m_{\mathcal{G}}$. Therefore $(\mathcal{G}, {}_{\mathcal{N},,\epsilon},)$ is flat by Proposition 5.5. In [10], F. Goodman and H. Wenzl showed that the fusion rule of $(\mathcal{G}, {}_{\mathcal{N},,\epsilon},)$ agrees with that of $SU(N)_L$-WZW model. We give another proof of their result in the next section.

Remark 6.1. (1) Strictly speaking, Wenzl deals with $w_{N,t,\epsilon}$ only when $\epsilon = -1$. However, it is clear that his arguments are applicable to the case $\epsilon = 1$. The results for $\mathfrak{A}(\mathfrak{w}_{\mathfrak{N},\mathfrak{t},\mathfrak{1}})$ also follows from those of $\mathfrak{A}(\mathfrak{w}_{\mathfrak{N},\mathfrak{t},-\mathfrak{1}})$, since the former is a 2-cocycle deformation of the latter (cf. [5]). Hence, ϵ may be viewed as a gauge parameter.
(2) In order to avoid using square roots of complex numbers, we use a different normalization of $w_{N,t,-1}$ from Wenzl [45]. For each $\mathbf{p} \in \mathcal{G}$ $(m \geq 1)$, we define $\kappa(\mathbf{p}) \in \mathbb{C}$ by

$$\kappa(\mathbf{p} \cdot \mathbf{q}) = \kappa(\mathbf{p})\kappa(\mathbf{q}) \quad (\mathbf{p} \in \mathcal{G}, \mathbf{q} \in \mathcal{G}^{\setminus}, , \setminus >), \tag{6.11}$$

$$\kappa(\lambda \,|\, i) = \prod_{k=i+1}^{N} A_{d(0 \,|\, i,k)+1}A_{d(0 \,|\, i,k)+2} \cdots A_{d(\lambda \,|\, i,k)} \quad ((\lambda \,|\, i) \in \mathcal{G}),$$

where $A_d = a_d/\sqrt{a_d a_{-d}}$ and $a_d = [d+1]/[2][d]$. Note that $\kappa(\mathbf{p})$ satisfies

$$\frac{\kappa(\mathbf{p})}{\kappa(\mathbf{p}^\dagger)} = A_{d(\mathbf{p})} \tag{6.12}$$

for each $\mathbf{p} \in \mathcal{G}[\searrow]$. By replacing the basis $\{\mathbf{p}\}$ of $\mathbb{C}\mathcal{G}$ with $\{\kappa(\mathbf{p})\mathbf{p}\}$, we obtain Wenzl's original expression of the Hecke algebra representation. It is also useful to use $\{\kappa(\mathbf{p})^{\mathbf{2}}\mathbf{p}\}$ instead of $\{\mathbf{p}\}$ (see §12). The corresponding Boltzmann weight $w^{\Sigma}_{N,t,\epsilon}$ satisfies

$$w^{\Sigma}_{N,t,\epsilon}\left[\mathbf{r}\overset{\mathbf{p}}{\underset{\mathbf{s}}{}}\mathbf{q}\right] := \left(\frac{\kappa(\mathbf{p}\cdot\mathbf{q})}{\kappa(\mathbf{r}\cdot\mathbf{s})}\right)^2 w_{N,t,\epsilon}\left[\mathbf{r}\overset{\mathbf{p}}{\underset{\mathbf{s}}{}}\mathbf{q}\right] = w_{N,t,\epsilon}\left[\mathbf{p}\overset{\mathbf{r}}{\underset{\mathbf{q}}{}}\mathbf{s}\right]. \tag{6.13}$$

We call $\{\mathbf{p}\}$ and $\{\kappa(\mathbf{p})^{\mathbf{2}}\mathbf{p}\}$ *rational basis of type* Ω and *type* Σ respectively.

7. THE ALGEBRA $\mathfrak{S}(A_{N-1};t)_\epsilon$

Applying Proposition 5.1 to $(\mathcal{G}, {}_{\mathcal{N},,\epsilon,\zeta})$, we obtain a CQT $\mathcal{V}$-face algebra $\mathfrak{A}(w_{N,t,\epsilon,\zeta})$ $= \mathrm{Cost}(\mathrm{w}_{\mathrm{N,t},\epsilon}, 0)$. In order to define the "(quantum) determinant" of $\mathfrak{A}(w_{N,t,\epsilon})$, we introduce an algebra $\Omega = \Omega_{N,L,\epsilon}$, which is a face-analogue of the exterior algebra. It is defined by generators $\omega(\mathbf{p})$ $(\mathbf{p} \in \mathcal{G};\ \geq\ )$ with defining relations:

$$\sum_{k\in\mathcal{V}} \omega(k) \ =\ 1, \tag{7.1}$$

$$\omega(\mathbf{p})\omega(\mathbf{q}) \ =\ \delta_{\mathfrak{r}(\mathbf{p})\mathfrak{s}(\mathbf{q})}\,\omega(\mathbf{p}\cdot\mathbf{q}), \tag{7.2}$$

$$\omega(\mathbf{p}) = -\epsilon\,\omega(\mathbf{p}^\dagger) \quad (\mathbf{p} \in \mathcal{G}[\searrow]), \tag{7.3}$$

$$\omega(\mathbf{p}) = \mathbf{0} \quad (\mathbf{p} \in \mathcal{G}[\rightarrow]). \tag{7.4}$$

It is easy to verify that $\Omega^m := \sum_{\mathbf{p}\in\mathcal{G}} \mathbb{C}\omega(\mathbf{p})$ becomes an $\mathfrak{A}(w_{N,t,\epsilon})$-comodule via

$$\omega(\mathbf{q}) \mapsto \sum_{\mathbf{p}\in\mathcal{G}} \omega(\mathbf{p}) \otimes \mathbf{e}\binom{\mathbf{p}}{\mathbf{q}} \quad (\mathbf{q} \in \mathcal{G}) \tag{7.5}$$

for each $m \geq 0$.

For each $m \geq 0$, we set

$$B\Omega^m = \big\{(\lambda,\, \lambda + \sum_{k\in I} \hat{i}\,) \in \mathcal{V} \,\big|\, \mathcal{I} \subset \{,\ldots,\mathcal{N}\},\,\mathrm{card(I) = m}\big\}. \tag{7.6}$$

Also we define $\mathcal{L}\colon \mathcal{G} \to \mathbb{Z}_{\geq\not{\prime}}$ by

$$\mathcal{L}(\lambda\,|\,,\ldots,\,) = \mathrm{Card}\{(\mathrm{k},\mathrm{l})|1 \leq \mathrm{k} < \mathrm{l} \leq \mathrm{N}, \mathrm{i_k} < \mathrm{i_l}\} \tag{7.7}$$

Proposition 7.1. *For each* $(\lambda,\mu) \in B\Omega^m$, $\mathcal{G}_{\lambda\mu} \neq \emptyset$ *and* $\omega_m(\lambda,\mu) := (-\epsilon)^{\mathcal{L}(\mathbf{p})}\omega(\mathbf{p})$ *does not depend on the choice of* $\mathbf{p} \in \mathcal{G}_{\lambda\mu}$. *Moreover* $\{\omega_m(\lambda,\mu)|(\lambda,\mu) \in B\Omega^m\}$ *is a basis of* Ω^m. *In particular,* $\Omega^m = 0$ *if* $m > N$.

Proof. We will prove this lemma by means of Bergman's diamond lemma [2], or rather its obvious generalization to the quotient algebras of $\mathbb{C}\langle\mathcal{G}\rangle$, where $\langle\mathcal{G}\rangle = \coprod \mathcal{G}$. We define a "reduction system" $S = S_1 \coprod S_2 \subset \langle\mathcal{G}\rangle \times \mathbb{C}\langle\mathcal{G}\rangle$ by setting

$$S_1 = \big\{(\mathbf{p}, -\epsilon\,\mathbf{p}^\dagger) \,\big|\, \mathbf{p} = (\lambda\,|\,\mathbf{i},\mathbf{j}) \in \mathcal{G}[\searrow],\ <|\big\},$$
$$S_2 = \big\{(\mathbf{p},\mathbf{0}) \,\big|\, \mathbf{p} = (\lambda\,|\,\mathbf{i_1},\ldots,\mathbf{i_m}) \in \mathcal{G},\ \geq,\, \mathrm{card}\{\mathrm{i_1},\ldots,\mathrm{i_m}\} < \mathrm{m}\big\}.$$

It is straightforward to verify that the quotient $\mathbb{C}\langle\mathcal{G}\rangle/\langle\mathcal{W} - \{\,|\,(\mathcal{W},\{) \in \mathcal{S}\rangle$ is isomorphic to Ω and that all ambiguities of S are resolvable. Next, we introduce a semigroup partial order $\leq$ on $\langle\mathcal{G}\rangle$ by setting $(\lambda\,|\,i_1,\ldots,i_m) < (\lambda\,|\,j_1,\ldots,j_n)$ if either $m < n$, or $m = n$ and $i_1 = j_1, \ldots, i_{k-1} = j_{k-1}$, $i_k > j_k$ for some $1 \leq k \leq m$. Then $\leq$ is compatible with S and satisfies the descending chain condition. This completes the proof of the proposition. □

For each $0 \leq m \leq N$, we set $\Lambda_m = \hat{1} + \cdots + \hat{m}$. As an immediate consequence of (5.22) and the the proposition above, we obtain the following result.

Proposition 7.2. *For each* $0 \leq m \leq N$*, we have* $\Omega^m \cong L_{(\Lambda_m,m)}$ *as* $\mathfrak{A}(w_{N,t,\epsilon})$*-comodules.*

Now we define the "determinant" $\det = \sum_{\lambda,\mu\in\mathcal{V}} \det\binom{\lambda}{\mu}$ of $\mathfrak{A}(w_{N,t,\epsilon})$ to be the group-like element which corresponds to the group-like comodule Ω^N and its basis $\{\bar{\omega}(\lambda)\}$ via Lemma 3.2, where $\bar{\omega}(\lambda) = D(\lambda)\omega_N(\lambda,\lambda)$ and

$$D(\lambda) = \prod_{1\leq i<j\leq N} \frac{[d(\lambda \,|\, i,j)]}{[d(0 \,|\, i,j)]} \quad (\lambda \in \mathcal{V}). \tag{7.8}$$

Explicitly, we have

$$\det\begin{pmatrix}\lambda\\ \mu\end{pmatrix} = \frac{D(\mu)}{D(\lambda)} \sum_{\mathbf{p}\in\mathcal{G}^N_{\lambda\lambda}} (-\epsilon)^{\mathcal{L}(\mathbf{p})+\mathcal{L}(\mathbf{q})} e\begin{pmatrix}\mathbf{p}\\ \mathbf{q}\end{pmatrix}, \tag{7.9}$$

where $\mathbf{q}$ denotes an arbitrary element of $\mathcal{G}^N_{\mu\mu}$. By (2.38) and (2.39) for $g = \det$, the quotient

$$\mathfrak{S}(A_{N-1};t)_\epsilon := \mathfrak{A}(w_{N,t,\epsilon})/(\det -1) \tag{7.10}$$

naturally becomes a $\mathcal{V}$-face algebra, which we call *$SU(N)_L$-SOS algebra.*
The proof of the following lemma will be given in §8 and §13.

Lemma 7.3. *For each* $\mathbf{p} \in \mathcal{G}$*, we have*

$$c_{\Omega^N\Omega^1}\left(\bar{\omega}(\mathfrak{s}(\mathbf{p})) \otimes \omega(\mathbf{p})\right) = \epsilon^{N-1}\zeta^{-N} t\omega(\mathbf{p}) \otimes \bar{\omega}(\mathfrak{r}(\mathbf{p})), \tag{7.11}$$

$$c_{\Omega^1\Omega^N}\left(\omega(\mathbf{p}) \otimes \bar{\omega}(\mathfrak{r}(\mathbf{p}))\right) = \epsilon^{N-1}\zeta^{-N} t\bar{\omega}(\mathfrak{s}(\mathbf{p})) \otimes \omega(\mathbf{p}). \tag{7.12}$$

Proposition 7.4. *The element* det *belongs to the center of* $\mathfrak{A}(w_{N,t,\epsilon})$*. Moreover, if* ζ *satisfies*

$$\zeta^N = \epsilon^{N-1}t, \tag{7.13}$$

then

$$\mathcal{R}^{\pm}(\det -1, a) = 0 = \mathcal{R}^{\pm}(a, \det -1) \quad (a \in \mathfrak{A}(w_{N,t,\epsilon,\zeta})). \tag{7.14}$$

Hence, $\mathfrak{S}(A_{N-1};t)_\epsilon$ *naturally becomes a quotient CQT* $\mathcal{V}$*-face algebra of* $\mathfrak{A}(w_{N,t,\epsilon,\zeta})$*.*

Proof. (cf. [13]) By computing the coaction of $\mathfrak{A}(w_{N,t,\epsilon})$ on $\omega(\mathbf{p}) \otimes \bar{\omega}(\mathfrak{r}(\mathbf{p}))$, in two ways via (7.11), we obtain the first assertion. We show the first equality of (7.14) for $\pm = +$ and $a = e\binom{\mathbf{p}}{\mathbf{q}}$ ($\mathbf{p},\mathbf{q} \in \mathcal{G}$, $\geq$). By (2.25) and (2.27), it suffices to show

$$\mathcal{R}^+\left(\det\begin{pmatrix}\mathfrak{r}(\mathbf{p})\\ \mathfrak{s}(\mathbf{p})\end{pmatrix}, e\begin{pmatrix}\mathbf{p}\\ \mathbf{q}\end{pmatrix}\right) = \delta_{\mathbf{pq}} \quad (\mathbf{p},\mathbf{q} \in \mathcal{G}). \tag{7.15}$$

For $m = 0$, this follows from (2.27) and (2.25). By computing the left-hand side of (7.11) via (3.5), we obtain (7.15) for $m = 1$. For $m \geq 2$, (7.15) follows from (2.23) and (2.38) for $g = \det$ by induction on m. □

Since the braiding of $\mathfrak{S}(A_{N-1};t)_\epsilon$ depends on the choice of the discrete parameter ζ satisfying (7.13), we sometimes write $\mathfrak{S}(A_{N-1};t)_{\epsilon,\zeta}$ instead of $\mathfrak{S}(A_{N-1};t)_\epsilon$.

To state our first main result, we recall the *fusion rule of $SU(N)_L$-WZWmodel* in conformal field theory. By [9], it is characterized as the structure constant $N^\nu_{\lambda\mu}$ of a commutative $\mathbb{Z}$-algebra $\mathcal{F}$ (called the *fusion algebra of $SU(N)_L$-WZWmodel*) with free basis $\{\chi_\lambda\}_{\lambda\in\mathcal{V}}$ (i.e., $\chi_\lambda\chi_\mu = \sum_{\nu\in\mathcal{V}} N^\nu_{\lambda\mu}\chi_\nu$) such that

$$N^\mu_{\lambda\Lambda_m} = \begin{cases} 1 & (\lambda,\mu) \in B\Omega^m \\ 0 & \text{otherwise} \end{cases} \tag{7.16}$$

for each $\lambda,\mu \in \mathcal{V}$ and $0 \leq m < N$. See Kac [25] or Walton [44] for an explicit formula of $N^\nu_{\lambda\mu}$.

Theorem 7.5 ([11]). (1) *For each* $\lambda \in \mathcal{V}$*, up to isomorphism there exists a unique right* $\mathfrak{S}(A_{N-1};t)_\epsilon$*-comodule* L_λ *such that*

$$\dim L_\lambda(0,\mu) = \delta_{\lambda\mu} \quad (\mu \in \mathcal{V}). \tag{7.17}$$

Moreover, we have

$$\mathfrak{S}(A_{N-1};t)_\epsilon \cong \bigoplus_{\lambda \in \mathcal{V}} \mathrm{End(L_\lambda)^*} \tag{7.18}$$

as coalgebras. In particular, L_λ *is irreducible for each* $\lambda \in \mathcal{V}$

(2) *The corepresentation ring* $K_0(\mathbf{Com}_{\mathfrak{S}(\mathbf{A_{N-1};t})_\epsilon})$ *is identified with the fusion algebra* $\mathcal{F}$ *of* $SU(N)_L$*-WZW model via* $\chi_\lambda = [L_\lambda]$*. That is, we have*

$$[L_0] = 1, \tag{7.19}$$

$$[L_\lambda][L_\mu] = \sum_{\nu \in \mathcal{V}} N^{\nu}_{\lambda\mu}[L_\nu]. \tag{7.20}$$

(3) *We have*

$$\dim(L_\nu(\lambda,\mu)) = N^{\mu}_{\lambda\nu}. \tag{7.21}$$

Proof. It is easy to verify that $(\lambda, m) \in \mathbf{\Lambda}_{\mathcal{G}_{\mathcal{N},\mathcal{L}}}$ if and only if $m \in |\lambda| + N\mathbb{Z}_{\geq\nvdash}$. Since $\mathbb{C}\mathcal{V}\det^{\backslash}\bar{\otimes}\mathcal{L}_{(\lambda,)} \cong L_{(\lambda,m+Nn)}$ by (3.2) and (5.18), we see that det satisfies all conditions of Lemma 3.3, where $\bar{\mathbf{\Lambda}} = \mathcal{V}$ and $\varphi(\lambda,n) = (\lambda, |\lambda| + Nn)$. Therefore, we have Part (1), (7.19) and

$$[L_\lambda][L_\mu] = \sum_{\nu \in \mathcal{V}} N^{\nu}_{\lambda\mu}(|\mu|)[L_\nu], \quad \dim(L_\nu(\lambda,\mu)) = N^{\mu}_{\lambda\nu}(m) \quad ((\nu,m) \in \mathbf{\Lambda}_{\mathcal{G}_{\mathcal{N},\mathcal{L}}}). \tag{7.22}$$

In particular, $\tilde{N}^{\nu}_{\lambda\mu} := N^{\nu}_{\lambda\mu}(m)$ does not depend on the choice of m. Using Proposition 7.2, (5.22) and (7.17), we obtain

$$[L_\lambda][L_{\Lambda_m}] = \sum_{\mu \in \mathcal{V}} \dim(L_\lambda \bar{\otimes} \Omega^m)(0,\mu)[L_\mu] = \sum_{\mu \in \mathcal{V}} \dim(\Omega^m(\lambda,\mu))[L_\mu]. \tag{7.23}$$

Thus the numbers $\tilde{N}^{\nu}_{\lambda\mu}$ satisfies the condition of $N^{\nu}_{\lambda\mu}$ stated above. □

Proposition 7.6. *The element* det *is not a zero-divisor of* $\mathfrak{A}(w_{N,t,\epsilon})$*. In particular, we have* $\det\binom{\lambda}{\mu} \neq 0$ *for each* $\lambda, \mu \in \mathcal{V}$*. Moreover, we have*

$$\mathrm{GLE}(\mathfrak{A}(\mathrm{w_{N,t,\epsilon}})) = \Big\{ \sum_{\lambda,\mu \in \mathcal{V}} \frac{\mathrm{c}(\lambda)}{\mathrm{c}(\mu)} \mathring{\mathrm{e}}_\lambda \mathrm{e}_\mu \mathrm{det^m} \;\Big|\; \mathrm{m} \in \mathbb{Z}_{\geq\nvdash},\ (\lambda) \in \mathbb{C}^\times\ (\lambda \in \mathcal{V}) \Big\}. \tag{7.24}$$

Proof. The first assertion follows from the fact that det is simply reducible (see the proof of the theorem above). By Theorem 5.3 (1), every group-like comodule of $\mathfrak{A}(w_{N,t,\epsilon})$ is isomorphic to $(\Omega^N)^{\overline{\otimes} m}$ for some $m \geq 0$. Hence the second assertion follows from Lemma 3.2. □

8. Module structure of Ω

Let $\mathfrak{H}$ be a CQT $\mathcal{V}$-face algebra over $\mathbb{K}$. As in case $\mathfrak{H}$ is a CQT bialgebra, the correspondence $a \mapsto \mathcal{R}^+(\ ,a)$ defines an antialgebra-coalgebra map from $\mathfrak{H}$ into the dual face algebra $\mathfrak{H}^\circ$ (cf. [16]). Let W be a right $\mathfrak{H}$-comodule. Combining the above map with the left action (3.11) of $\mathfrak{H}^*$ on W, we obtain a right action of $\mathfrak{H}$ on W given by

$$wa = \sum_{(w)} w_{(0)} \mathcal{R}^+(w_{(1)}, a) \quad (w \in W, a \in \mathfrak{H}). \tag{8.1}$$

Let V be another $\mathfrak{H}$-comodule. Then we have

$$(v \otimes w)\, a = \sum_{(a)} va_{(1)} \otimes wa_{(2)} \quad (v \in V(\lambda, \nu), w \in W(\nu, \mu), a \in \mathfrak{H}). \tag{8.2}$$

If $\mathfrak{H}$ has an antipode and W is finite-dimensional, then we have

$$\langle va, w \rangle = \langle v, wS^{-1}(a) \rangle \quad (v \in W^{\vee}, w \in W), \tag{8.3}$$

by (3.6) and (2.29).

In this section, we give an explicit description of the right $\mathfrak{A}(w_{N,t,\epsilon})$-module structure of Ω. By (7.5) and (5.5), we obtain the following.

Lemma 8.1. *For each* $\mathbf{s} \in \mathcal{G}$ *and* $\mathbf{p}, \mathbf{q} \in \mathcal{G}^{\setminus}$ $(n \geq 0)$, *we have*

$$\omega(\mathbf{s})\, \mathbf{e}\binom{\mathbf{p}}{\mathbf{q}} = \sum_{\mathbf{r} \in \mathcal{G}} \mathbf{w}_{\mathbf{N},\mathbf{t},\epsilon} \left[\mathbf{p} \overset{\mathbf{s}}{\underset{\mathbf{r}}{}} \mathbf{q}\right] \omega(\mathbf{r}) \quad (\mathbf{s} \in \mathcal{G}, \mathbf{p}, \mathbf{q} \in \mathcal{G}^{\setminus}, \setminus \geq). \tag{8.4}$$

In particular, we have

$$\Omega(\lambda, \mu) e\binom{\mathbf{p}}{\mathbf{q}} \in \delta_{\lambda, \mathfrak{s}(\mathbf{p})} \delta_{\mu, \mathfrak{s}(\mathbf{q})}\, \Omega(\mathfrak{r}(\mathbf{p}), \mathfrak{r}(\mathbf{q})). \tag{8.5}$$

Lemma 8.2. *Let* (λ, μ) *be an element of* $B\Omega^m$ $(m > 0)$ *and* $\mathbf{p} = (\lambda \,|\, i_1, \ldots, i_m)$ *an element of* $\mathcal{G}_{\lambda\mu}$. *Define the set* I *and* $C(\lambda \,|\, k, l) \in \mathbb{C}$ $(k \neq l)$ *by* $I = \{i_1, \ldots, i_m\}$ *and*

$$C(\lambda \,|\, k, l) = \frac{[d(\lambda \,|\, k, l) + 1]}{[d(\lambda \,|\, k, l)]} \tag{8.6}$$

respectively. Then for each $(\lambda \,|\, i), (\mu \,|\, j) \in \mathcal{G}$, *we have*:

$$\omega(\mathbf{p})\mathbf{e}\binom{\lambda \,|\, \mathbf{i}}{\mu \,|\, \mathbf{j}} \;=\; (-\zeta)^{-\mathbf{m}} \mathbf{t}^{-\mathbf{d}(\lambda |\, \mathbf{i},\mathbf{j})} \frac{\mathbf{1}}{[\mathbf{d}(\lambda |\, \mathbf{i}, \mathbf{j})]} \prod_{\mathbf{k} \in \mathbf{I} \setminus \{\mathbf{i}\}} \mathbf{C}(\lambda \,|\, \mathbf{i}, \mathbf{k})\, \omega(\lambda + \hat{\mathbf{i}} \,|\, \mathbf{i_2}, \ldots, \mathbf{i_m}, \mathbf{j}) \qquad (i = i_1, j \notin I), \tag{8.7}$$

$$\omega(\mathbf{p})\mathbf{e}\binom{\lambda \,|\, \mathbf{i}}{\mu \,|\, \mathbf{j}} \;=\; -\delta_{\mathbf{ij}} (-\zeta)^{-\mathbf{m}} \mathbf{t} \prod_{\mathbf{k} \in \mathbf{I} \setminus \{\mathbf{i}\}} \mathbf{C}(\lambda \,|\, \mathbf{i}, \mathbf{k})\, \omega_{\mathbf{m}}(\lambda + \hat{\mathbf{i}} \,|\, \mathbf{i_2}, \ldots, \mathbf{i_m}, \mathbf{i}) \qquad (i = i_1, j \in I), \tag{8.8}$$

$$\omega_m(\mathbf{p})\mathbf{e}\binom{\lambda \,|\, \mathbf{i}}{\mu \,|\, \mathbf{j}} \;=\; \delta_{\mathbf{ij}} (\epsilon \zeta^{-\mathbf{1}})^{\mathbf{m}} \prod_{\mathbf{k} \in \mathbf{I}} \mathbf{C}(\lambda \,|\, \mathbf{i}, \mathbf{k})\, \omega_{\mathbf{m}}(\lambda + \hat{\mathbf{i}} \,|\, \mathbf{i_1}, \ldots, \mathbf{i_m}) \qquad (i, j \notin I), \tag{8.9}$$

$$\omega_m(\mathbf{p})\mathbf{e}\binom{\lambda \,|\, \mathbf{i}}{\mu \,|\, \mathbf{j}} \;=\; \mathbf{0} \qquad (\mathbf{i} \notin \mathbf{I}, \mathbf{j} \in \mathbf{I}). \tag{8.10}$$

Proof. These formulas are proved by induction on m in a similar manner. Here we give the proof of (8.7). Since Ω^m is a quotient module of $\mathbb{C}\mathcal{G}^{-} \bar{\otimes} \mathbb{C}\mathcal{G}$, the left-hand side of (8.7) is rewritten as $\sum_q A_q B_q$ with

$$A_q = \omega_{m-1}(\lambda, \nu) e\binom{\lambda \,|\, i}{\nu \,|\, q}, \quad B_q = \omega_1(\nu, \mu) e\binom{\nu \,|\, q}{\mu \,|\, j} \tag{8.11}$$

by (8.2), where $\nu = \lambda + \hat{i}_1 + \cdots + \hat{i}_{m-1}$ and the summation is taken over for all $1 \leq q \leq N$ such that $(\nu \,|\, q) \in \mathcal{G}$. Since $(\nu + \hat{q}, \mu + \hat{j}) \in \mathcal{G}$ only if $q = i_m$ or j, we have

$$\omega_m(\lambda, \mu) e\binom{\lambda \,|\, i}{\mu \,|\, j} = \begin{cases} A_{i_m} B_{i_m} + A_j B_j & (\nu \,|\, j) \in \mathcal{G} \\ A_{i_m} B_{i_m} & \text{otherwise.} \end{cases} \tag{8.12}$$

Using inductive assumption, we see that the right-hand side of (8.12) equals

$$(-\zeta)^{-m}t^{-d(i,j)}\left(\frac{1}{[d(i,i_m)][d(i_m,j)]}+\frac{[d(i_m,j)-1]}{[d(i,j)][d(i_m,j)]}\right)$$
$$\prod_{n=2}^{m-1}C(\lambda\,|\,i,i_n)\,\omega_m(\lambda+\hat{i},\mu+\hat{j}) \tag{8.13}$$

if $(\nu\,|\,j)\in\mathcal{G}$, where $d(k,l)=d(\lambda\,|\,k,l)$. Applying $[a+b+1]+[a][b]=[a+1][b+1]$ to $a=d(i,i_m)$ and $b=d(i_m,j)-1$, we see that (8.13) equals the right-hand side of (8.7) (even if $(\nu\,|\,j)\notin\mathcal{G}$). Next suppose that $(\nu\,|\,j)\notin\mathcal{G}$. It suffices to verify that the second term in the parentheses of (8.13) is zero. In case $j=1$, we obtain $\nu_1=L$. Using this together with $1\notin I$, we see that $\lambda_1=L$. On the other hand, since $(\mu\,|\,1)\in\mathcal{G}$, we have $L-1\geq\mu_1=L-\delta_{i_mN}$. Hence, $i_m=N$ and $d(i_m,j)-1=-L-N$. In case $j>1$, we obtain $d(i_m,j)-1=0$ in a similar manner. Thus we complete the proof of (8.7). □

The following lemma is frequently used in the sequel.

Lemma 8.3. *As a right $\mathfrak{A}(w_{N,t,\epsilon})$-module, $\mathbb{C}\mathcal{G}\cong\Omega$ is irreducible. Hence $\mathbb{C}\mathcal{G}$ is also irreducible as a left $\mathfrak{A}(w_{N,t,\epsilon})^*$-module.*

Proof. Let W be a non-zero submodule of $\mathbb{C}\mathcal{G}$. Since $W=\bigoplus_{\lambda\mu}W\mathring{e}_\lambda e_\mu$ and $W\mathring{e}_\lambda e_\mu\subset\mathbb{C}\mathcal{G}_{\lambda\mu}$, we have $\mathbf{s_0}\in\mathbf{W}$ for some $\mathbf{s_0}=(\lambda\,|\,\mathbf{i})\in\mathcal{G}$. To show $\mathbb{C}\mathcal{G}=\mathbf{s_0}\,\mathfrak{A}(\mathbf{w_{N,t,\epsilon}})$, we introduce the oriented graph $\mathcal{H}$ determined by $\mathcal{H}=\mathcal{G}$ and

$$\operatorname{card}(\mathcal{H}_{\mathbf{sr}})=\begin{cases}1 & \mathbf{s}\,\mathbf{e}\binom{\mathbf{p}}{\mathbf{q}}\in\mathbb{C}^\times\mathbf{r}\quad(\exists\mathbf{p},\mathbf{q}\in\mathcal{G})\\ 0 & \text{otherwise.}\end{cases} \tag{8.14}$$

It suffices to show that $\langle\mathcal{H}\rangle_{\mathbf{s_0\,r}}\neq\emptyset$ for every $\mathbf{r}\in\mathcal{G}$, where $\langle\mathcal{H}\rangle_{\mathbf{s\,r}}=\cup_m\mathcal{H}_{\mathbf{s\,r}}$. We note that

$$\mathcal{H}_{(\mu\,|\,|)\,(\mu+\hat{\imath}\,|\,)}\neq\emptyset\quad\text{if }\ (\mu\,|\,\mathrm{j}),(\mu+\hat{\mathrm{j}}\,|\,\mathrm{k})\in\mathcal{G}\text{ and }\mathrm{j}\neq\mathrm{k}, \tag{8.15}$$

$$\mathcal{H}_{(\mu\,|\,|)\,(\mu+\hat{}\,|\,|)}\neq\emptyset\quad\text{if }\ (\mu\,|\,\mathrm{j}),(\mu+\hat{\mathrm{k}}\,|\,\mathrm{j})\in\mathcal{G}\text{ and }\mathrm{j}\neq\mathrm{k}, \tag{8.16}$$

$$\mathcal{H}_{(\mu\,|\,|)\,(\mu+\hat{\imath}\,|\,|)}\neq\emptyset\quad\text{if }\ (\mu\,|\,\mathrm{j},\mathrm{j})\in\mathcal{G}. \tag{8.17}$$

Using (8.16), we obtain $\langle\mathcal{H}\rangle_{\mathbf{s_0\,s_1}}\neq\emptyset$, where $\mathbf{s_1}=(L\Lambda_{i-1}+\sum_{k=i}^{N-1}\lambda_k\hat{k}\,|\,i)$. Suppose $i\neq N$. Using (8.17), and then using (8.16), we obtain $\langle\mathcal{H}\rangle_{\mathbf{s_1\,s_2}}\neq\emptyset$, where $\mathbf{s_2}=((L-1)\Lambda_{N-1}+\Lambda_{i-1}\,|\,i)$. Using (8.15) and (8.16) respectively, we obtain $\langle\mathcal{H}\rangle_{\mathbf{s_2\,s_3}}$, $\langle\mathcal{H}\rangle_{\mathbf{s_4\,(0|1)}}\neq\emptyset$ and $\langle\mathcal{H}\rangle_{\mathbf{s_3\,s_4}}\neq\emptyset$ respectively, where $\mathbf{s_3}=((L-1)\Lambda_{N-1}+\Lambda_{N-2}\,|\,N-1)$. and $\mathbf{s_4}=(\Lambda_{N-2}\,|\,N-1)$. Therefore we obtain $\langle\mathcal{H}\rangle_{\mathbf{s_0(0|1)}}\neq\emptyset$ if $i\neq N$. By similar consideration, we also obtain $\langle\mathcal{H}\rangle_{\mathbf{s_0(0|1)}}\neq\emptyset$ in case $i=N$, and also, $\langle\mathcal{H}\rangle_{(|)\mathbf{r}}\neq\emptyset$ for every $\mathbf{r}\in\mathcal{G}$. Thus, we have verified the first assertion. The second assertion is obvious since the image of $a\mapsto\mathcal{R}^+(\ ,a)$ is a subalgebra of $\mathfrak{A}(w_{N,t,\epsilon})^*$. □

Now we begin to prove Lemma 7.3. By (8.5) and Lemma 8.2, we have

$$\bar{\omega}(\nu)e\begin{pmatrix}\lambda\,|\,i\\ \mu\,|\,j\end{pmatrix}\ =\ \epsilon^{N-1}\zeta^{-N}t\delta_{ij}\delta_{\nu\lambda}\delta_{\nu\mu}\bar{\omega}(\nu+\hat{i}). \tag{8.18}$$

Using (8.2) and this equality, we see that both $\Omega^N\bar{\otimes}\mathbb{C}\mathcal{G}\to\mathbb{C}\mathcal{G}$; $\bar{\omega}(\mathfrak{s}(\mathbf{p}))\otimes\mathbf{p}\mapsto\mathbf{p}\otimes\bar{\omega}(\mathfrak{r}(\mathbf{p}))$ and $\mathbb{C}\mathcal{G}\bar{\otimes}\Omega^N\to\mathbb{C}\mathcal{G}$; $\mathbf{p}\otimes\bar{\omega}(\mathfrak{s}(\mathbf{p}))\mapsto\mathbf{p}\otimes\bar{\omega}(\mathfrak{r}(\mathbf{p}))$ are isomorphisms of right $\mathfrak{A}(w_{N,t,\epsilon})$-modules. Hence, by Lemma 8.3 and Schur's Lemma, we have

$$c_{\Omega^N\Omega^1}(\bar{\omega}(\mathfrak{s}(\mathbf{p}))\otimes\mathbf{p})=\vartheta\,\mathbf{p}\otimes\bar{\omega}(\mathfrak{r}(\mathbf{p}))\quad(\mathbf{p}\in\mathcal{G}) \tag{8.19}$$

for some constant ϑ. We will prove $\vartheta=\epsilon^{N-1}\zeta^{-N}t$ in §12.

9. TRANSPOSES AND COMPLEX CONJUGATES

The following proposition is an immediate consequence of the following reflection symmetry:

$$w_{N,t,\epsilon}\left[\mathbf{r}\overset{\mathbf{p}}{\underset{\mathbf{s}}{\,}}\mathbf{q}\right] = \left(\frac{\kappa(\mathbf{r}\cdot\mathbf{s})}{\kappa(\mathbf{p}\cdot\mathbf{q})}\right)^2 w_{N,t,\epsilon}\left[\mathbf{p}\overset{\mathbf{r}}{\underset{\mathbf{q}}{\,}}\mathbf{s}\right], \tag{9.1}$$

where κ is as in (6.11).

Proposition 9.1. *There exists an algebra-anticoalgebra map* $\mathfrak{A}(w_{N,t,\epsilon}) \to \mathfrak{A}(w_{N,t,\epsilon})$; $a \mapsto a^{\mathsf{T}}$ *given by*

$$e\begin{pmatrix}\mathbf{p}\\ \mathbf{q}\end{pmatrix}^{\mathsf{T}} = \left(\frac{\kappa(\mathbf{p})}{\kappa(\mathbf{q})}\right)^2 e\begin{pmatrix}\mathbf{q}\\ \mathbf{p}\end{pmatrix} \quad (\mathbf{p},\mathbf{q}\in\mathcal{G},\ \geq\ ). \tag{9.2}$$

Moreover it satisfies $(a^{\mathsf{T}})^{\mathsf{T}} = a$ *and*

$$\mathcal{R}^{\pm}\,(a^{\mathsf{T}},\, b^{\mathsf{T}}) = \mathcal{R}^{\pm}\,(b,\, a) \tag{9.3}$$

for each $a, b \in \mathfrak{A}(w_{N,t,\epsilon})$ *and* $\zeta\in\mathbb{C}^{\times}$.

The following proposition is needed to construct the "cofactor matrix."

Proposition 9.2. *The element* det *satisfies* $\det^{\mathsf{T}} = \det$. *Hence,* T *induces an algebra-anticoalgebra involution of* $\mathfrak{S}(A_{N-1};t)_\epsilon$, *which satisfies* (9.3).

Proof. Since $\det^{\mathsf{T}}$ is a group-like element of $\mathfrak{A}(w_{N,t,\epsilon})$, we have

$$\det{}^{\mathsf{T}} = g\det; \quad g = \sum_{\lambda,\mu\in\mathcal{V}} \frac{c(\lambda)}{c(\mu)}\,\overset{\circ}{e}_\lambda e_\mu \tag{9.4}$$

by (7.24), where $c(\lambda)$ $(\lambda\in\mathcal{V})$ denotes some nonzero constant. Since both det and $\det^{\mathsf{T}}$ are central and det is not a zero divisor, g is central. Hence, by Lemma 8.3 and Schur's lemma, we have $\mathbf{p}\,\mathbf{g} = \mathbf{c}\,\mathbf{p}$ $(\mathbf{p}\in\mathcal{G})$ for some $c\in\mathbb{C}$. Hence we have $c(\lambda) = c^{|\lambda|}c(0)$ for each $\lambda\in\mathcal{V}$. In order to prove $c = 1$, we compute $\det\binom{0}{\hat{1}}^{\mathsf{T}}$ in two ways. Using

$$[d(\mathbf{p}) - \mathbf{1}]\,\mathbf{e}\begin{pmatrix}\mathbf{p}\\ \mathbf{q}\end{pmatrix} = -\epsilon[\mathbf{d}(\mathbf{p}) + \mathbf{1}]\,\mathbf{e}\begin{pmatrix}\mathbf{p}^{\dagger}\\ \mathbf{q}\end{pmatrix} \quad (\mathbf{p}\in\ \mathcal{G}^{\mathbf{2}}[\searrow],\, \mathbf{q}\in\mathcal{G}^{\mathbf{2}}[\downarrow]), \tag{9.5}$$

we obtain

$$\sum_{i=1}^{k}(-\epsilon)^{\mathcal{L}(\mathbf{p_i})} e\begin{pmatrix}\mathbf{p_i}\\ \mathbf{0}\,|\,\mathbf{1},\ldots,\mathbf{N}\end{pmatrix} = (-\epsilon)^{\mathcal{L}(\mathbf{p_k})}[k]^2\, e\begin{pmatrix}\mathbf{p_k}\\ \mathbf{0}\,|\,\mathbf{1},\ldots,\mathbf{N}\end{pmatrix} \tag{9.6}$$

by induction on k, where $\mathbf{p_i} = (\hat{\mathbf{1}}\,|\,\mathbf{2},\ldots,\mathbf{i},\mathbf{1},\mathbf{i}+\mathbf{1},\ldots,\mathbf{N})$. Substituting $k = N$ in this equality, we get

$$\begin{aligned}\det\begin{pmatrix}0\\ \hat{1}\end{pmatrix}^{\mathsf{T}} &= c^{|\hat{1}|-|0|}\det\begin{pmatrix}\hat{1}\\ 0\end{pmatrix}\\ &= (-\epsilon)^{N(N-1)/2+\mathcal{L}(\mathbf{p_N})}c[N]\,e\begin{pmatrix}\hat{1}\,|& 2,& 3,& \ldots,& N,& 1\\ 0\,|& 1,& 2,& \ldots,& N-1,& N\end{pmatrix}.\end{aligned}$$

On the other hand, using (7.9) and (9.2), we see that the right-hand side of the above equality agrees with $c\det\binom{0}{\hat{1}}^{\mathsf{T}}$. This completes the proof of the proposition. □

Next suppose $t = \exp(\pm\frac{\pi i}{N+L})$, or $-\exp(\pm\frac{\pi i}{N+L})$ with $N + L \in 2\mathbb{Z}$. Then, we have $\kappa(\mathbf{p}) > \mathbf{0}$ for each $\mathbf{p}\in\mathcal{G}$ $(m > 0)$. Moreover, for each ζ with $|\zeta| = 1$, the Boltzmann weight $w_{N,t}$ satisfies

$$w_{N,t,\epsilon}^{-1}\left[\mathbf{r}\overset{\mathbf{p}}{\underset{\mathbf{s}}{\,}}\mathbf{q}\right] = \left(\frac{\kappa(\mathbf{r}\cdot\mathbf{s})}{\kappa(\mathbf{p}\cdot\mathbf{q})}\right)^2 \overline{w_{N,t,\epsilon}\left[\mathbf{p}\overset{\mathbf{r}}{\underset{\mathbf{q}}{\,}}\mathbf{s}\right]}. \tag{9.7}$$

Similarly to Proposition 9.1 and Proposition 9.2, we obtain the following.

Proposition 9.3. *Set* $t = \pm\exp(\pm\frac{\pi i}{N+L})$ *if* $N + L \in 2\mathbb{Z}$*, and* $t = \exp(\pm\frac{\pi i}{N+L})$ *if* $N + L \in 1 + 2\mathbb{Z}$*. Then for each solution* ζ *of* (7.13)*, both* $\mathfrak{S}(A_{N-1}; t)_{\epsilon,\zeta}$ *and* $\mathfrak{A}(w_{N,t,\epsilon,\zeta})$ *are compact CQT* $\mathcal{V}$*-face algebras of unitary type with costar structure*

$$e\binom{\mathbf{p}}{\mathbf{q}}^{\times} = \left(\frac{\kappa(\mathbf{p})}{\kappa(\mathbf{q})}\right)^2 e\binom{\mathbf{q}}{\mathbf{p}} \quad (\mathbf{p}, \mathbf{q} \in \mathcal{G},\ \geq\ ). \tag{9.8}$$

In fact, both $\mathfrak{S}(A_{N-1}; t)_\epsilon$ and $\mathfrak{A}(w_{N,t,\epsilon})$ are spanned by unitary matrix corepresentations $[e_u\binom{\mathbf{p}}{\mathbf{q}}]_{\mathbf{p},\mathbf{q}\in\mathcal{G}}$ $(m \geq 0)$ given by

$$e_u\binom{\mathbf{p}}{\mathbf{q}} = \frac{\kappa(\mathbf{q})}{\kappa(\mathbf{p})}\, e\binom{\mathbf{p}}{\mathbf{q}} \quad (\mathbf{p}, \mathbf{q} \in \mathcal{G},\ \geq\ ). \tag{9.9}$$

10. ANTIPODES AND RIBBON FUNCTIONALS

Lemma 10.1. *The* $\mathcal{V}$*-face algebra* $\mathfrak{S}(A_{N-1}; t)_\epsilon$ *has an antipode given by*

$$S\left(e\binom{\lambda\,|\,i}{\mu\,|\,j}\right) = (-\epsilon)^{i+j}\frac{D(\lambda+\hat{i})}{D(\mu+\hat{j})}\sum_{\mathbf{a}}(-\epsilon)^{\mathcal{L}(\mathbf{a})+\mathcal{L}(\mathbf{b})} e\binom{\mathbf{a}}{\mathbf{b}}, \tag{10.1}$$

where $\mathbf{b}$ *denotes an arbitrary element of* $\mathcal{G}^{\mathcal{N}-}_{\lambda+\hat{\ }\lambda}$ *and the summation is taken over all* $\mathbf{a} \in \mathcal{G}^{\mathcal{N}-}_{\mu+\hat{\imath}\,\mu}$*. Moreover, we have*

$$S^2\left(e\binom{\mathbf{p}}{\mathbf{q}}\right) = \frac{D(\mathfrak{r}(\mathbf{p}))\mathbf{D}(\mathfrak{s}(\mathbf{q}))}{\mathbf{D}(\mathfrak{s}(\mathbf{p}))\mathbf{D}(\mathfrak{r}(\mathbf{q}))}\, e\binom{\mathbf{p}}{\mathbf{q}} \quad (\mathbf{p}, \mathbf{q} \in \mathcal{G},\ \geq\ ). \tag{10.2}$$

Proof. (cf. [38], [12]) Let $Y_{(\lambda\,|\,i)\,(\mu\,|\,j)}$ denote the right-hand side of (10.1) viewed as an element of $\mathfrak{A}(w_{N,t,\epsilon})$. By [17, §7], it suffices to verify that

$$\sum_{\mathbf{r}\in\mathcal{G}} Y_{\mathbf{pr}}X_{\mathbf{rq}} = \delta_{\mathbf{pq}}e_{\mathfrak{r}(\mathbf{p})}\det, \quad \sum_{\mathbf{r}\in\mathcal{G}} X_{\mathbf{pr}}Y_{\mathbf{rq}} = \delta_{\mathbf{pq}}\overset{\circ}{e}_{\mathfrak{s}(\mathbf{p})}\det, \tag{10.3}$$

where $X_{\mathbf{pq}} = e\binom{\mathbf{p}}{\mathbf{q}}$ $(\mathbf{p}, \mathbf{q} \in \mathcal{G})$. We define a basis $\{\bar{\omega}(\tilde{\mathbf{p}}) \mid \mathbf{p} \in \mathcal{G}\}$ of Ω^{N-1} by $\bar{\omega}(\lambda+\hat{i}, \lambda) = (-\epsilon)^{i-1}D(\lambda+\hat{i})\omega_{N-1}(\lambda+\hat{i}, \lambda)$, where $\tilde{\mathbf{p}} = (\mu, \lambda) \in B\Omega^{N-1}$ for $\mathbf{p} = (\lambda, \mu) \in \mathcal{G}$. Then, the coaction of $\mathfrak{A}(w_{N,t,\epsilon})$ on Ω^{N-1} is given by $\bar{\omega}(\tilde{\mathbf{q}}) \mapsto \sum_{\mathbf{p}\in\mathcal{G}} \bar{\omega}(\tilde{\mathbf{p}}) \otimes Y_{\mathbf{qp}}$ and the multiplication of Ω gives maps

$$\Omega^{N-1}\bar{\otimes}\Omega^1 \to \Omega^N; \quad \bar{\omega}(\tilde{\mathbf{q}}) \otimes \omega(\mathbf{p}) \mapsto \delta_{\mathbf{pq}}\bar{\omega}(\mathfrak{r}(\mathbf{p})),$$

$$\Omega^1\bar{\otimes}\Omega^{N-1} \to \Omega^N; \quad \omega(\mathbf{p}) \otimes \bar{\omega}(\tilde{\mathbf{q}}) \mapsto \delta_{\mathbf{pq}}(-\epsilon)^{\mathbf{N-1}}\frac{\mathbf{D}(\mathfrak{r}(\mathbf{p}))}{\mathbf{D}(\mathfrak{s}(\mathbf{p}))}\bar{\omega}(\mathfrak{s}(\mathbf{p})).$$

Since these maps are compatible with the coaction of $\mathfrak{A}(w_{N,t,\epsilon})$, we have the first formula of (10.3) and

$$\sum_{\mathbf{r}} W_{\mathbf{rp}}Y_{\mathbf{qr}} = \delta_{\mathbf{pq}}e_{\mathfrak{s}(\mathbf{p})}\det, \tag{10.4}$$

where $W_{\mathbf{pq}} \in \mathfrak{A}(w_{N,t,\epsilon})$ denotes the right-hand side of (10.2). Applying $\intercal$ to this equality, we obtain

$$\sum_{\mathbf{r}} X_{\mathbf{pr}}Z_{\mathbf{rq}} = \delta_{\mathbf{pq}}\overset{\circ}{e}_{\mathfrak{s}(\mathbf{p})}\det; \quad Z_{\mathbf{pq}} = \frac{D(\mathfrak{r}(\mathbf{p}))\mathbf{D}(\mathfrak{s}(\mathbf{q}))\kappa(\mathbf{p})^{\mathbf{2}}}{\mathbf{D}(\mathfrak{s}(\mathbf{p}))\mathbf{D}(\mathfrak{r}(\mathbf{q}))\kappa(\mathbf{q})^{\mathbf{2}}}\, Y^{\intercal}_{\mathbf{qp}}. \tag{10.5}$$

Computing $\sum_{\mathbf{rs}} Y_{\mathbf{pr}}X_{\mathbf{rs}}Z_{\mathbf{sq}}$ in two ways, we obtain $Y_{\mathbf{pq}} = Z_{\mathbf{pq}}$. This proves the second equality of (10.3). Finally, Computing $\sum_{\mathbf{rs}} W_{\mathbf{rq}}S(X_{\mathbf{sr}})S^2(X_{\mathbf{ps}})$ in the algebra $\mathfrak{S}(A_{N-1}; t)_\epsilon$ in two ways, we obtain (10.2). □

Proposition 10.2. *For each t and ζ, $\mathfrak{S}(A_{N-1};t)_{\epsilon,\zeta}$ becomes a coribbon Hopf $\mathcal{V}$-face algebra, whose braiding, antipode S and modified ribbon functional $\mathcal{M} = \mathcal{M}_\infty$ are given by* (5.5), (10.1) *and the following formulas*:

$$\mathcal{M}\left(\rceil\binom{\mathbf{p}}{\mathbf{q}}\right) = \delta_{\mathbf{pq}}\frac{\mathcal{D}(\mathfrak{r}(\mathbf{p}))}{\mathbf{D}(\mathfrak{s}(\mathbf{p}))} \quad (\mathbf{p},\mathbf{q}\in\mathcal{G},\ \geq\ ). \tag{10.6}$$

Moreover, we have

$$\dim_q(L_\lambda) = D(\lambda) \tag{10.7}$$

for each $\lambda\in\mathcal{V}$. When N is even, there exists another ribbon functional $\mathcal{M}_{-\infty}$ given by

$$\mathcal{M}_{-\infty}\left(\rceil\binom{\mathbf{p}}{\mathbf{q}}\right) = \delta_{\mathbf{pq}}\,(-\infty)^{|\mathfrak{r}(\mathbf{p})|-|\mathfrak{s}(\mathbf{p})|}\frac{\mathcal{D}(\mathfrak{r}(\mathbf{p}))}{\mathbf{D}(\mathfrak{s}(\mathbf{p}))} \quad (\mathbf{p},\mathbf{q}\in\mathcal{G},\ \geq\ ). \tag{10.8}$$

The quantum dimension of the corresponding ribbon category is given by

$$\dim_q^{-1}(L_\lambda) = (-1)^{|\lambda|}D(\lambda). \tag{10.9}$$

Proof. Let $\mathcal{M}\in\mathfrak{H}^*$ be as in (10.6). Using (10.2), we see that $\mathcal{M}$ satisfies (2.43). Hence, it suffices to verify that

$$(\mathcal{U}_\infty\mathcal{U}_\in)\left(\rceil\binom{\mathbf{p}}{\mathbf{q}}\right) = \mathcal{M}^{\in}\left(\rceil\binom{\mathbf{p}}{\mathbf{q}}\right) \quad (\mathbf{p},\mathbf{q}\in\mathcal{G}). \tag{10.10}$$

for each $m\geq 0$. By (2.34) and (2.43), $\mathcal{U}_\infty\mathcal{M}^{-\infty}$ is a central element of $\mathfrak{A}(w_{N,t,\epsilon})^*$. Hence by Lemma 8.3 and Schur's lemma, we have $\mathcal{U}_\infty\mathbf{p} = \vartheta\mathcal{M}\mathbf{p}$ ($\mathbf{p}\in\mathcal{G}$) for some constant ϑ. Using (2.31), (2.29) and (10.2), we compute

$$\mathcal{U}_\infty^{-\infty}\left(\rceil\binom{\prime\,|\,\infty}{\prime\,|\,\infty}\right) = \sum_{\mathbf{r}\in\mathcal{G}}\mathcal{R}^+\left(S^2\left(e\binom{\mathbf{r}}{\mathbf{0}\,|\,\mathbf{1}}\right), e\binom{0\,|\,1}{\mathbf{r}}\right) \tag{10.11}$$

$$= \sum_{\nu=2\hat{1},\hat{1}+\hat{2}}\frac{D(0)D(\nu)}{D(\hat{1})^2}w_{N,t}\begin{bmatrix}0 & \hat{1}\\ \hat{1} & \nu\end{bmatrix} \tag{10.12}$$

$$= \zeta^{-1}t^N\frac{1}{[N]}. \tag{10.13}$$

This shows that $\vartheta = \zeta t^{-N}$, and similarly, we obtain $\mathcal{U}_\in\mathbf{p} = \vartheta^{-\mathbf{1}}\mathcal{M}\mathbf{p}$ ($\mathbf{p}\in\mathcal{G}$). This proves (10.10) for $m=1$. For $m\geq 2$, (10.10) follows from (10.10) for $m=1$ by induction on m, using the fact that both $\mathcal{M}^{\in}$ and $\mathcal{U}_\infty\mathcal{U}_\in$ are group-like (cf. (2.32), (2.33)). The second assertion follows from (10.6) and Lemma 3.1. □

We denote by $\mathfrak{S}(A_{N-1};t)^{\iota}_{\epsilon,\zeta}$ the coribbon Hopf $\mathcal{V}$-face algebra $(\mathfrak{S}(A_{N-1};t)_{\epsilon,\zeta},\mathcal{M}_\iota)$, where $\iota=\pm 1$ if N is even and $\iota = 1$ if N is odd.

Lemma 10.3. *If $t = \exp(\pm\frac{\pi i}{N+L})$, or $t = -\exp(\pm\frac{\pi i}{N+L})$ and N is odd, then we have $D(\lambda) > 0$ for every $\lambda\in\mathcal{V}$. If $t = -\exp(\pm\frac{\pi i}{N+L})$ and N is even, then we have $(-1)^{|\lambda|}D(\lambda) > 0$.*

Proof. Straightforward. □

Proposition 10.4. *When $t = \exp(\pm\frac{\pi i}{N+L})$, or $t = -\exp(\pm\frac{\pi i}{N+L})$ and $N, L\in 1+2\mathbb{Z}$, the Woronowicz functional $\mathcal{Q}$ of $\mathfrak{S}(A_{N-1};t)_\epsilon$ is given by* (10.6)*. While when $t = -\exp(\pm\frac{\pi i}{N+L})$ and $N, L\in 2\mathbb{Z}$, $\mathcal{Q}$ is given by* (10.8)*.*

Proof. We will prove the first assertion. We set

$$M = \left[\mathcal{M}\left(\rceil_\sqcap\binom{\mathbf{p}}{\mathbf{q}}\right)\right]_{\mathbf{p},\mathbf{q}\in\mathcal{G}} \qquad Q = \left[\mathcal{Q}\left(\rceil_\sqcap\binom{\mathbf{p}}{\mathbf{q}}\right)\right]_{\mathbf{p},\mathbf{q}\in\mathcal{G}} \tag{10.14}$$

where $\mathcal{Q}$ denotes the Woronowicz functional of $\mathfrak{S}(A_{N-1};t)_\epsilon$ and $e_u\binom{\mathbf{p}}{\mathbf{q}}$ is as in (9.9). By (2.43), (4.6) and Lemma 8.3, we have $M = \vartheta Q$ for some ϑ. Since M is positive by (10.6) and the lemma above, we have $\vartheta > 0$. Since the quantum dimension satisfies $\dim_q L = \dim_q L^\vee$ for every L, we have $\mathrm{Tr(M)} = \mathrm{Tr(M^{-1})}$. By (4.7), this proves $M = Q$. Now the assertion $\mathcal{M}\big(\rceil\sqcap\binom{\mathbf{p}}{\mathbf{q}}\big) = \mathcal{Q}\big(\rceil\sqcap\binom{\mathbf{p}}{\mathbf{q}}\big)$ $(\mathbf{p},\mathbf{q} \in \mathcal{G})$ easily follows from the fact that both $\mathcal{M}$ and $\mathcal{Q}$ are group-like, by induction on m. □

11. The modular tensor category

Let $\mathcal{C}$ be a ribbon category, which is additive over a field $\mathbb{K}$. We say that $\mathcal{C}$ is *semisimple* if there exist a set $\mathcal{V}_{\mathcal{C}}$, an involution ${}^\vee : \mathcal{V}_{\mathcal{C}} \to \mathcal{V}_{\mathcal{C}}$, an element $0 \in \mathcal{V}_{\mathcal{C}}$ and simple objects $L^{\mathcal{C}}_\lambda$ $(\lambda \in \mathcal{V}_{\mathcal{C}})$ such that every object of $\mathcal{C}$ is isomorphic to a finite direct sum of $L^{\mathcal{C}}_\lambda$'s, and that

$$L^{\mathcal{C}}_0 \cong \mathbf{1}, \quad (\mathbf{L}^{\mathcal{C}}_\lambda)^\vee \cong \mathbf{L}^{\mathcal{C}}_{\lambda^\vee}, \tag{11.1}$$

$$\mathcal{C}(\mathcal{L}^{\mathcal{C}}_\lambda, \mathcal{L}^{\mathcal{C}}_\mu) = \begin{cases} \mathbb{K} & (\lambda = \mu) \\ 0 & (\lambda \neq \mu) \end{cases} \tag{11.2}$$

for each $\lambda, \mu \in \mathcal{V}^{\mathcal{C}}$, where $\mathbf{1}$ denotes the unit object of $\mathcal{C}$. For a semisimple ribbon category $\mathcal{C}$, we define its *fusion rule* $N^\nu_{\lambda\mu}$ $(\lambda, \mu, \nu \in \mathcal{V}_{\mathcal{C}})$ and *S-matrix* $S^{\mathcal{C}} = [S^{\mathcal{C}}_{\lambda\mu}]_{\lambda,\mu\in\mathcal{V}_{\mathcal{C}}}$ by

$$[L^{\mathcal{C}}_\lambda][L^{\mathcal{C}}_\mu] = \sum_{\nu\in\mathcal{V}_{\mathcal{C}}} N^\nu_{\lambda,\mu}[L^{\mathcal{C}}_\nu], \tag{11.3}$$

$$S^{\mathcal{C}}_{\lambda\mu} = \mathrm{Tr_q}(\mathrm{c}_{\mathrm{L}^{\mathcal{C}}_\mu \mathrm{L}^{\mathcal{C}}_\lambda} \circ \mathrm{c}_{\mathrm{L}^{\mathcal{C}}_\lambda \mathrm{L}^{\mathcal{C}}_\mu}). \tag{11.4}$$

By definition, we have

$$S^{\mathcal{C}}_{\lambda 0} = \dim_q L^{\mathcal{C}}_\lambda. \tag{11.5}$$

Since the twist θ satisfies

$$c_{WV} \circ c_{VW} = \theta_{V\otimes W} \circ (\theta_V \otimes \theta_W)^{-1}, \tag{11.6}$$

$S^{\mathcal{C}}$ satisfies

$$S^{\mathcal{C}}_{\lambda\mu} = \sum_{\nu\in\mathcal{V}_{\mathcal{C}}} \frac{\theta_\nu}{\theta_\lambda\theta_\mu} N^\nu_{\lambda\mu} \dim_q(L^{\mathcal{C}}_\nu), \tag{11.7}$$

where $\theta_\lambda \in \mathbb{K}$ is defined by $\theta_{L^{\mathcal{C}}_\lambda} = \theta_\lambda \mathrm{id}_{\mathrm{L}^{\mathcal{C}}_\lambda}$. Moreover, $S = S^{\mathcal{C}}$ satisfies the following *Verlinde's formula* (cf. [43, 32, 40]):

$$S_{\nu,0} \sum_{\xi\in\mathcal{V}} N^\xi_{\lambda\mu} S_{\xi\nu} = S_{\lambda\nu} S_{\mu\nu}, \tag{11.8}$$

where λ, μ and ν denote arbitrary elements of $\mathcal{V} = \mathcal{V}_{\mathcal{C}}$. Let $\mathcal{C}$ be a semisimple ribbon category. We say that $\mathcal{C}$ is a *modular tensor category* (or *MTC*) if $\mathcal{V}_{\mathcal{C}}$ is finite and the matrix $S^{\mathcal{C}}$ is invertible. If, in addition, $\mathcal{C}$ is unitary as a ribbon category, then it is called a *unitary MTC*. It is known that each (unitary) MTC gives rise to a (unitary) 3-dimensional topological quantum field theory (TQFT), hence, in particular, an invariant of 3-manifolds of Witten-Reshetikhin-Turaev type (cf. V. Turaev [40]).

The most well-known example of MTC is obtained as a certain semisimple quotient $\mathcal{C}(\mathfrak{g}, \kappa)$ of a category of representations of the quantized enveloping algebra $U_q(\mathfrak{g})$ of finite type in the case when q is a root of unity [1, 8, 28, 41]. When $\mathfrak{g} = \mathfrak{sl}_{\mathfrak{N}}$, the simple objects L^U_λ of $\mathcal{C}(\mathfrak{sl}_{\mathfrak{N}}, \mathfrak{N} + \mathfrak{L})$ $(L \geq 1)$ are also indexed by the set $\mathcal{V} = \mathcal{V}_{\mathcal{NL}}$ given by (6.1) and the fusion rules agree with those of $SU(N)_L$-WZW model. The quantum dimension and the constant θ_λ for $\mathcal{C}(\mathfrak{sl}_{\mathfrak{N}}, \mathfrak{N} + \mathfrak{L})$ are given by

$$\dim_q(L^U_\lambda) = D(\lambda)_{t_0}, \quad \theta_\lambda = \zeta_0^{(\lambda|\lambda+2\rho)^\sim} \tag{11.9}$$

respectively, where $\zeta_0 = \exp(\frac{\pi i}{N(N+L)})$, $t_0 = \zeta_0^N$, $\rho = \Lambda_1 + \cdots + \Lambda_{N-1}$, $(\,|\,)^\sim = N(\,|\,)$ and $(\,|\,)$ denotes the usual inner product of $\mathbb{R}^{\mathbb{N}}$. Moreover, the S-matrix of $\mathcal{C}(\mathfrak{sl}_{\mathfrak{N}}, \mathfrak{N}+\mathfrak{L})$ is given by $S^U = S^1(\zeta_0)$. Here, for each primitive $2N(N+L)$-th root ζ of unity, we define the matrix $S^\iota(\zeta)$ by the following Kac-Peterson formula (cf. [28]):

$$S^\iota(\zeta)_{\lambda\mu} = \iota^{|\lambda|+|\mu|} \frac{\sum_{w\in\mathfrak{S}_{\mathfrak{N}}}(-1)^{l(w)}\zeta^{-2(w(\lambda+\rho)\,|\,\mu+\rho)^\sim}}{\sum_{w\in\mathfrak{S}_{\mathfrak{N}}}(-1)^{l(w)}\zeta^{-2(w(\rho)\,|\,\rho)^\sim}}, \tag{11.10}$$

where $\iota = \pm 1$ if $N \in 2\mathbb{Z}$, $\iota = 1$ if $N \in 1+2\mathbb{Z}$ and the action of the symmetric group $\mathfrak{S}_{\mathfrak{N}}$ on $\sum_i \mathbb{C}\hat{\beth}$ is given by $w\hat{i} = \widehat{w(i)}$. Note that $(\lambda\,|\,\mu)^\sim \in \mathbb{Z}$ for every $\lambda, \mu \in \bigoplus_i \mathbb{Z}\Lambda_\beth$.

Lemma 11.1. *Let ζ be a primitive $2N(N+L)$-th root of unity and $t = \zeta^N$. Then the matrix $S = S^\iota(\zeta)$ is both symmetric and invertible, and satisfies Verlinde's formula* (11.8)*. Moreover, we have:*

$$S^\iota(\zeta)_{\lambda 0} = \iota^{|\lambda|} D(\lambda)_t, \tag{11.11}$$

$$S^\iota(\zeta)_{\Lambda_q\Lambda_r} = \iota^{q+r}\sum_s (\zeta t)^{-2qr} t^{2s(q+r-s+1)} D(\Lambda_{q+r-s}+\Lambda_s)_t. \tag{11.12}$$

for each $\lambda \in \mathcal{V}_{\mathcal{NL}}$ and $0 < r \le q < N$, where the summation in (11.12) *is taken over* $\max\{0, q+r-N\} \le s \le r$*.*

Proof. Suppose $\zeta = \zeta_0$. Then the formula (11.12) follows from (11.7) for $\mathcal{C}(\mathfrak{sl}_{\mathfrak{N}}, \mathfrak{N}+\mathfrak{L})$, (7.16) and (11.9). In this case, the other assertions also follow from the results for $\mathcal{C}(\mathfrak{sl}_{\mathfrak{N}}, \mathfrak{N}+\mathfrak{L})$. For other ζ, the assertions follow from Galois theory for $\mathbb{Q}(\zeta_{\nvdash})/\mathbb{Q}$. $\square$

Lemma 11.2. *Let $N^\nu_{\lambda\mu}$ be the fusion rules of $SU(N)_L$-WZW models and let S and S' be symmetric matrices whose entries are indexed by $\mathcal{V} = \mathcal{V}_{\mathcal{NL}}$. If these satisfy Verlinde's formula* (11.8)*, $S_{\lambda 0} = S'_{\lambda 0} \neq 0$ $(\lambda \in \mathcal{V})$ and*

$$S_{\Lambda_q\Lambda_r} = S'_{\Lambda_q\Lambda_r} \quad (0 < r \le q < N), \tag{11.13}$$

then we have $S = S'$.

Proof. We recall that there exists an algebra surjection from $\mathbb{Z}[\curvearrowleft_{\nVdash}, \ldots, \curvearrowleft_{\mathbb{N}}]^{\mathfrak{S}_{\mathfrak{N}}}$ onto $\mathcal{F}$ (cf. Theorem 7.5 (2)), which sends the Schur function $s_{(\lambda_1,\ldots,\lambda_N)}$ (see e.g. [31]) to $[L_\lambda]$ for each $\lambda \in \mathcal{V}$ (see e.g. [9]). For each $\xi = (\xi_1, \ldots, \xi_m)$ such that $N > \xi_1 \ge \ldots \ge \xi_m > 0$, we define $E_\xi \in \mathcal{F}$ to be the image of the elementary symmetric function e_ξ via this map, that is $E_\xi = [\Omega^{\xi_1}]\cdots[\Omega^{\xi_m}]$. Since $\{e_\xi\}$ is a basis of $\mathbb{Z}[\curvearrowleft_{\nVdash}, \ldots, \curvearrowleft_{\mathbb{N}}]^{\mathfrak{S}_{\mathfrak{N}}}$, $\{E_\xi\}$ spans $\mathcal{F}$. We define the symmetric bilinear forms $\mathcal{S}$ and $\mathcal{S}'$ on $\mathcal{F}$ by setting

$$\mathcal{S}([\mathcal{L}_\lambda], [\mathcal{L}_\mu]) = \mathcal{S}_{\lambda\mu}, \quad \mathcal{S}'([\mathcal{L}_\lambda], [\mathcal{L}_\mu]) = \mathcal{S}'_{\lambda\mu}. \tag{11.14}$$

Then, Verlinde's formula for S is rewritten as

$$\mathcal{S}\left(\dashv\lfloor, \frac{[\mathcal{L}_\nu]}{\mathcal{S}_{\nu\prime}}\right) = \mathcal{S}\left(\dashv, \frac{[\mathcal{L}_\nu]}{\mathcal{S}_{\nu\prime}}\right)\mathcal{S}\left(\lfloor, \frac{[\mathcal{L}_\nu]}{\mathcal{S}_{\nu\prime}}\right), \tag{11.15}$$

where $\nu \in \mathcal{V}$ and $a, b \in \mathcal{F}$. By (11.13) and this formula, we obtain

$$\mathcal{S}(\mathcal{E}_\xi, [\Omega^\nabla]) = \mathcal{S}'(\mathcal{E}_\xi, [\Omega^\nabla]) \tag{11.16}$$

for each ξ and $0 \le r < N$, or equivalently, we obtain

$$\mathcal{S}([\mathcal{L}_\lambda], [\Omega^\nabla]) = \mathcal{S}'([\mathcal{L}_\lambda], [\Omega^\nabla]) \tag{11.17}$$

for each $\lambda \in \mathcal{V}$ and $0 \leq r < N$. Repeating similar consideration, we conclude that $S_{\lambda\mu} = S'_{\lambda\mu}$ holds for every $\lambda, \mu \in \mathcal{V}$. □

For $\mathfrak{S} = \mathfrak{S}(\mathfrak{A}_{\mathfrak{N}-\mathfrak{1}}; \mathfrak{t})^{\iota}_{\epsilon,\zeta}$, we denote the semisimple ribbon category $\mathbf{Com}^{\mathbf{f}}_{\mathfrak{S}}$ (resp. unitary ribbon category $\mathbf{Com}^{\mathbf{fu}}_{\mathfrak{S}}$) by $\mathcal{C}_{\mathfrak{S}}(\mathcal{A}_{\mathcal{N}-\infty}, \sqcup)^{\iota}_{\epsilon,\zeta}$ (resp. $\mathcal{C}^{\sqcap}_{\mathfrak{S}}(\mathcal{A}_{\mathcal{N}-\infty}, \sqcup)_{\epsilon,\zeta}$).

Theorem 11.3. *Let $N \geq 2$ and $L \geq 1$ be integers, $\iota = \pm 1$ if $N \in 2\mathbb{Z}$, $\iota = 1$ if $N \in 1 + 2\mathbb{Z}$ and $\epsilon = \pm 1$. Let ζ be a primitive $2N(N+L)$-th root of unity.*
(1) *Suppose N is odd or $\epsilon = 1$ and set $t = \zeta^N$. Then the category $\mathcal{C}_{\mathfrak{S}}(\mathcal{A}_{\mathcal{N}-\infty}, \sqcup)^{\iota}_{\epsilon,\zeta}$ is a modular tensor category with S-matrix $S^{\iota}(\zeta)$.*
(2) *Suppose N is even, $\epsilon = -1$ and $t := -\zeta^N$ is a primitive $2(N+L)$-th root of unity. (Note that this implies $L \in 2\mathbb{Z}$). Then the category $\mathcal{C}_{\mathfrak{S}}(\mathcal{A}_{\mathcal{N}-\infty}, \sqcup)^{\iota}_{-\infty,\zeta}$ is a modular tensor category with S-matrix $S^{-\iota}(\zeta)$.*

Proof. We will prove Part (2). By (11.5), (10.7), (10.9), (11.11) and $D(\lambda)_{-t} = (-1)^{|\lambda|} D(\lambda)_t$, we have $S^{\mathfrak{S}}_{\lambda 0} = S^{-\iota}(\zeta)_{\lambda 0}$. Hence by the lamma above, it suffices to show that $S^{\mathfrak{S}}_{\Lambda_q \Lambda_r} = S^{-\iota}(\zeta)_{\Lambda_q \Lambda_r}$ for each $0 < r \leq q < N$. As we will see in the next section, the action of $c_{\Omega^r \Omega^q} \circ c_{\Omega^q \Omega^r}$ on the one-dimensional space $(\Omega^q \bar{\otimes} \Omega^r)(0, \Lambda_{q+r-s} + \Lambda_s) = \mathbb{C}\omega_{\shortparallel}(\nvdash, \Lambda_{\shortparallel}) \otimes \omega_{\diagdown}(\Lambda_{\shortparallel}, \Lambda_{\shortparallel+\diagdown-\sim} + \Lambda_{\sim})$ is given by the scalar $(\zeta t)^{-2qr} t^{2s(q+r-s+1)}$. Hence, by (3.14), we obtain

$$
\begin{aligned}
S^{\mathfrak{S}}_{\Lambda_q \Lambda_r} &= \sum_s \mathrm{Tr}\big(\mathcal{M}_{\iota} \circ \rfloor_{\Omega^{\triangledown}\Omega^{\amalg}} \circ \rfloor_{\Omega^{\amalg}\Omega^{\triangledown}}\big|_{(\Omega^{\amalg}\bar{\otimes}\Omega^{\triangledown})(\prime, \Lambda_{\amalg+\triangledown-\int} + \Lambda_{\int})}\big) \\
&= (-\iota)^{p+q} \sum_s (\zeta t')^{-2qr} t'^{2s(q+r-s+1)} D(\Lambda_{q+r-s} + \Lambda_s)_{t'},
\end{aligned}
\tag{11.18}
$$

where $t' = \zeta^N$ and the summation is taken over $\max\{0, q+r-N\} \leq s \leq r$. Since the right-hand side of (11.18) equals $S^{-\iota}(\zeta)_{\Lambda_q \Lambda_r}$ by (11.12), this completes the proof of Part (2). □

Corollary 11.4. *Let $N \geq 2$ and $L \geq 1$ be integers, $\epsilon = \pm 1$ and $t = \exp(\pm\frac{\pi i}{N+L})$. If $N + L \in 1 + 2\mathbb{Z}$, $\mathcal{C}^{\sqcap}_{\mathfrak{S}}(\mathcal{A}_{\mathcal{N}-\infty}, \sqcup)_{\epsilon,\zeta}$ is a unitary MTC provided that $\epsilon = 1$ or N is odd, where ζ denotes an arbitrary primitive $2N(N+L)$-th root of unity such that $\zeta^N = t$. If $N + L \in 2\mathbb{Z}$, $\mathcal{C}^{\sqcap}_{\mathfrak{S}}(\mathcal{A}_{\mathcal{N}-\infty}, \pm\sqcup)_{\epsilon,\zeta}$ is a unitary MTC for each primitive $2N(N+L)$-th root ζ of unity such that $\zeta^N = \pm\epsilon^{N-1} t$.*

Remark 11.1. (1) When $N \in 1 + 2\mathbb{Z}$, $\mathfrak{S}(\mathfrak{A}_{\mathfrak{N}-\mathfrak{1}}, \mathfrak{t})^{\iota}_{-1,\zeta}$ is isomorphic to a 2-cocycle deformation of $\mathfrak{S}(\mathfrak{A}_{\mathfrak{N}-\mathfrak{1}}, \mathfrak{t})^{\iota}_{1,\zeta}$. Hence $\mathcal{C}_{\mathfrak{S}}(\mathcal{A}_{\mathcal{N}-\infty}, \sqcup)^{\iota}_{\infty,\zeta}$ and $\mathcal{C}_{\mathfrak{S}}(\mathcal{A}_{\mathcal{N}-\infty}, \sqcup)^{\iota}_{-\infty,\zeta}$ are equivalent.

(2) For $\mathfrak{g} = \mathfrak{so}_{\mathfrak{N}}$ and $\mathfrak{sp}_{\mathfrak{N}}$, a category-theoretic construction of unitary MTC's related to $\mathcal{C}(\mathfrak{g}, \kappa)$ is given by Turaev and Wenzl [42].

12. BRAIDINGS ON Ω

In this section, we give some explicit calculation of the braiding $c_{q,r} := c_{\Omega^q \Omega^r}$ in order to complete the proof of Lemma 7.3 and Theorem 11.3. Since the braiding is a natural transformation and the multiplication of Ω gives a $\mathfrak{S}(A_{N-1}; t)_{\epsilon}$-comodule map $m_{q,r} \colon \Omega^q \bar{\otimes} \Omega^r \to \Omega^{q+r}$, $c_{q,r}$ satisfies

$$
c_{q+q',r} \circ (m_{q,q'} \bar{\otimes} \mathrm{id}_{\Omega^{\mathrm{r}}}) = (\mathrm{id}_{\Omega^{\mathrm{r}}} \bar{\otimes} \mathrm{m}_{\mathrm{q},\mathrm{q}'}) \circ (\mathrm{c}_{\mathrm{q},\mathrm{r}} \bar{\otimes} \mathrm{id}_{\Omega^{\mathrm{q}'}}) \circ (\mathrm{id}_{\Omega^{\mathrm{q}}} \bar{\otimes} \mathrm{c}_{\mathrm{q}',\mathrm{r}}),
\tag{12.1}
$$

$$
c_{q,r+r'} \circ (\mathrm{id}_{\Omega^{\mathrm{q}}} \bar{\otimes} \mathrm{m}_{\mathrm{r},\mathrm{r}'}) = (\mathrm{m}_{\mathrm{r},\mathrm{r}'} \bar{\otimes} \mathrm{id}_{\Omega^{\mathrm{q}}}) \circ (\mathrm{id}_{\Omega^{\mathrm{r}}} \bar{\otimes} \mathrm{c}_{\mathrm{q},\mathrm{r}'}) \circ (\mathrm{c}_{\mathrm{q},\mathrm{r}} \bar{\otimes} \mathrm{id}_{\Omega^{\mathrm{r}'}}).
\tag{12.2}
$$

Lemma 12.1. *For each* $1 \le p < N$ *and* $1 \le q \le N-p$, *we have*

$$\begin{aligned} &c_{q,1}\big(\omega\,(\Lambda_p\,|\,p+1,\dots,p+q)\otimes\omega\,(\Lambda_{p+q}\,|\,1)\big)\\ &\qquad = -(-\zeta)^{-q}t^{p+1}\frac{[q]}{[p+1]}\,\omega\,(\Lambda_p\,|\,p+1)\otimes\omega\,(\Lambda_{p+1}\,|\,p+2,\dots,p+q,1)\\ &\qquad + (-\zeta)^{-q}(-\epsilon)^q\frac{[p+q+1]}{[p+1]}\,\omega\,(\Lambda_p\,|\,1)\otimes\omega\,\big(\Lambda_p+\hat{1}\,|\,p+1,\dots,p+q\big)\,. \end{aligned} \tag{12.3}$$

Proof. Suppose (12.3) is valid for each p and for some $q < N-p$. Then, by (12.1), we obtain

$$c_{1+q,1}\big(\omega\,(\Lambda_p\,|\,p+1,\dots,p+q+1)\otimes\omega\,(\Lambda_{p+q+1}\,|\,1)\big) \tag{12.4}$$

$$\begin{aligned} &= \ (\mathrm{id}_{\Omega_1}\bar{\otimes}\mathrm{m}_{1,\mathrm{q}})\circ(\mathrm{c}_{1,1}\bar{\otimes}\mathrm{id}_{\Omega^\mathrm{q}})\Big[\omega(\Lambda_\mathrm{p}|\mathrm{p}+1)\otimes\\ &\quad\big\{-(-\zeta)^{-q}t^{p+2}\frac{[q]}{[p+2]}\,\omega\,(\Lambda_{p+1}\,|\,p+2)\otimes\omega\,(\Lambda_{p+2}\,|\,p+3,\dots,p+q+1,1)\\ &\quad+(-\zeta)^{-q}(-\epsilon)^q\frac{[p+q+2]}{[p+2]}\,\omega\,(\Lambda_{p+1}\,|\,1)\otimes\omega\,\big(\Lambda_{p+1}+\hat{1}\,|\,p+2,\dots,p+q+1\big)\big\}\Big]\\ &= \ \left\{-t^{p+2}\frac{[q]}{[p+2]}\,w_{N,t}\begin{bmatrix}\Lambda_p & \Lambda_{p+1}\\ \Lambda_{p+1} & \Lambda_{p+2}\end{bmatrix}+\frac{[p+q+2]}{[p+2]}w_{N,t}\begin{bmatrix}\Lambda_p & \Lambda_{p+1}\\ \Lambda_{p+1} & \Lambda_{p+1}+\hat{1}\end{bmatrix}\right\}\\ &\qquad(-\zeta)^{-q}\omega(\Lambda_p|p+1)\otimes\omega\,(\Lambda_{p+1}\,|\,p+2,\dots,p+q+1,1)\\ &\qquad\quad+(-\zeta)^{-q}(-\epsilon)^q\frac{[p+q+2]}{[p+2]}\,w_{N,t}\begin{bmatrix}\Lambda_p & \Lambda_{p+1}\\ \Lambda_p+\hat{1} & \Lambda_{p+1}+\hat{1}\end{bmatrix}\\ &\qquad\qquad\qquad\omega\,(\Lambda_p\,|\,1)\otimes\omega\,\big(\Lambda_p+\hat{1}\,|\,p+1,\dots,p+q+1\big)\,. \end{aligned}$$

Computing the right-hand side of the above equality, we obtain (12.3) for $q+1$. ☐

Using (12.3) for $p=1$, $q=N-1$ together with (12.1), we obtain

$$c_{N,1}\big(\omega\,(0|\,1,\dots,N)\otimes\omega\,(0\,|\,1)\big) = -(-\zeta)^{-N}t\,[N]\,\omega\,(0\,|\,1)\otimes\omega\,(1\,|\,2,\dots,N,1)\,. \tag{12.5}$$

This shows that the constant ϑ in (8.19) equals $\epsilon^{N-1}\zeta^{-N}t$ and completes the proof of Lemma 7.3.

Lemma 12.2. *We have the following relations*:

$$\begin{aligned} &c_{q,1}\Big(\omega\,(\Lambda_p\,|\,p+1,\dots,p+s,1,\dots,q-s)\otimes\omega\,(\Lambda_{p+s}+\Lambda_{q-s}\,|\,q-s+1)\Big)\\ &\in -(-\zeta)^{-q}t^{p-q+s+1}\frac{[s]}{[p+1]}\,\omega\,(\Lambda_p\,|\,p+1)\otimes\omega\,(\Lambda_{p+1}\,|\,p+2\dots,p+s,1,\dots,q-s+1)\\ &\qquad\qquad + \Omega^1\,(\Lambda_p\,|\,1)\otimes\Omega^q\,\big(\Lambda_p+\hat{1},\Lambda_{p+s}+\Lambda_{q-s+1}\big)\\ &\qquad\qquad\qquad (1 \le p < N, 1 \le s \le N-p, s < q \le p+2s-1), \end{aligned} \tag{12.6}$$

$$\begin{aligned} &c_{q,r}\Big(\omega\,(\Lambda_p\,|\,p+1,\dots,p+q)\otimes\omega\,(\Lambda_{p+q}\,|\,1,\dots,r)\Big)\\ &\qquad\in(-1)^r(-\zeta)^{-qr}t^{pr+r}\frac{[q]!\,[p]!}{[p+r]!\,[q-r]!}\,\omega\,(\Lambda_p\,|\,p+1,\dots,p+r)\\ &\qquad\qquad\otimes\omega\,(\Lambda_{p+r}\,|\,p+r+1,\dots,p+q,1,\dots,r)\\ &\qquad\qquad\qquad+\sum_{\lambda\neq\Lambda_{p+r}}\Omega^r\,(\Lambda_p,\lambda)\otimes\Omega^q\,(\lambda,\Lambda_{p+q}+\Lambda_r)\\ &\qquad\qquad\qquad (0 \le p \le N-1, 0 \le q \le N-p, 0 \le r \le q), \end{aligned} \tag{12.7}$$

where $[n]! = [n]\cdots[2][1]$ *and* $[0]! = 1$.

Proof. The relation (12.6) follows from (12.1), (12.3) and

$$c_{q-s,1}\big(\omega\,(\Lambda_{p+s}\,|\,1,\dots,q-s)\otimes\omega\,(\Lambda_{p+s}+\Lambda_{q-s}\,|\,q-s+1)\big)$$
$$=(-\zeta t)^{s-q}\omega\,(\Lambda_{p+s}\,|\,1)\otimes\omega\,\big(\Lambda_{p+s}+\hat{1}\,|\,2,\dots,q-s+1\big)\,. \tag{12.8}$$

The relation (12.7) is easily proved by induction on r, using (12.2) and (12.6). □

Using (12.7), (12.2) and

$$c_{q,r-s}\big(\omega\,(0\,|\,1,\dots,q)\otimes\omega\,(\Lambda_q\,|\,q+1,\dots,q+r-s)\big)$$
$$=(-\zeta t)^{qs-qr}\omega\,(0\,|\,1,\dots,r-s)\otimes\omega\,(\Lambda_{r-s}\,|\,r-s+1,\dots,q+r-s)\,, \tag{12.9}$$

we obtain the following.

Lemma 12.3. *For each* $0\le q,r<N$ *and* $\max\{0,q+r-N\}\le s\le\min\{q,r\}$, *we have*

$$c_{q,r}\big(\omega\,(0\,|\,1,\dots,q)\otimes\omega\,(\Lambda_q\,|\,q+1,\dots,q+r-s,1,\dots s)\big)$$
$$=\ (-1)^s(-\zeta t)^{-qr}t^{s(q+r-s+1)}\frac{[q]!\,[r-s]!}{[r]!\,[q-s]!}$$
$$\omega\,(0\,|\,1,\dots,r)\otimes\omega\,(\Lambda_r\,|\,r+1,\dots,q+r-s,1,\dots,s)\,. \tag{12.10}$$

As an immediate consequence of the lemma above, we see that $c_{r,q}\circ c_{q,r}$ acts on $(\Omega^q\bar{\otimes}\Omega^r)(0,\,\Lambda_{q+r-s}+\Lambda_s)$ as the scalar $(\zeta t)^{-2qr}t^{2s(q+r-s+1)}$. Thus we complete the proof of Theorem 11.3.

13. ABF MODELS AND $SU(2)_L$-SOS ALGEBRAS

In this section, we give an explicit description of the representation theory of $\mathfrak{S}(A_1;t)_\epsilon$. We identify $\mathcal{G}=\mathcal{G}_{,\mathcal{L}}$ with the Dynkin diagram of type A_{L+1}:



(13.1)

Also, we identify $\mathcal{V}$ and $\mathcal{G}_|$ with $\{0,1,\cdots,L\}$ and

$$\{\,(i_0,i_1,\cdots,i_k)|\;0\le i_0,\cdots,i_k\le L,\;|i_\nu-i_{\nu-1}|=1\,(1\le\nu\le k)\} \tag{13.2}$$

respectively. We define the set $\mathcal{B}$ by

$$\mathcal{B}=\left\{\begin{pmatrix}\|\\ \rangle|\end{pmatrix}\middle|\begin{array}{c}\rangle,|,\|\in\mathcal{V},\;|-\|\le\ \le\ +|,\\ \rangle+|+\|\in\in\mathbb{Z},\;\beth+\gimel+\daleth\le\nvDash\mathbb{L}\end{array}\right\}. \tag{13.3}$$

Then, we have

$$N_{ij}^k=\begin{cases}1 & \left(\binom{k}{ij}\in\mathcal{B}\right)\\ 0 & (\text{otherwise})\,.\end{cases} \tag{13.4}$$

In order to simplify the formula for quantum invariants stated in the introduction, we use the rational basis of type Σ instead of type Ω (cf. Remark 6.1). The corresponding Boltzmann weight $w=w^{\Sigma}_{2,t,\epsilon}$ is given by

$$w\begin{bmatrix}i & i\pm1\\ i\pm1 & i\end{bmatrix}=-\zeta^{-1}\,\frac{\pm t^{\mp(i+1)}}{[i+1]},\quad w\begin{bmatrix}i & i\pm1\\ i\mp1 & i\end{bmatrix}=\zeta^{-1}\epsilon\,\frac{[i+1\pm1]}{[i+1]}, \tag{13.5}$$

$$w\begin{bmatrix}i & i\pm1\\ i\pm1 & i\pm2\end{bmatrix}=\zeta^{-1}t,\quad w\Big[\text{otherwise}\Big]=0. \tag{13.6}$$

Next, we recall a realization of L_k introduced in [14]. Let Σ be an algebra generated by the symbols $\sigma(\mathbf{p})$ ($\mathbf{p} \in \mathcal{G},\ \geq$) with defining relations:

$$\sum_{k \in \mathcal{V}} \sigma(k) \;=\; 1, \tag{13.7}$$

$$\sigma(\mathbf{p})\sigma(\mathbf{q}) \;=\; \delta_{\mathfrak{r}(\mathbf{p})\mathfrak{s}(\mathbf{q})}\, \sigma(\mathbf{p} \cdot \mathbf{q}), \tag{13.8}$$

$$\sigma(i, i+1, i) = \epsilon\sigma(i, i-1, i) \quad (0 < i < L), \tag{13.9}$$

$$\sigma(0, 1, 0) = \sigma(L, L-1, L) = 0. \tag{13.10}$$

We define the grading $\Sigma = \bigoplus_{k \geq 0} \Sigma^k$ via $\Sigma^k = \operatorname{span}\{\sigma(\mathbf{p}) \mid \mathbf{p} \in \mathcal{G}\}$. Then each component Σ^k becomes a right $\mathfrak{S}(A_1; t)_\epsilon$-comodule via

$$\sigma(\mathbf{q}) \mapsto \sum_{\mathbf{p} \in \mathcal{G}} \sigma(\mathbf{p}) \otimes \mathbf{e}\binom{\mathbf{p}}{\mathbf{q}} \quad (\mathbf{q} \in \mathcal{G}). \tag{13.11}$$

For each $\binom{k}{ij} \in \mathcal{B}$, the element $\sigma_k(i,j) := \epsilon^{\mathcal{L}(\mathbf{q})}\sigma(\mathbf{q})$ does not depend on the choice of $\mathbf{q} \in \mathcal{G}_{|}$, where $\mathcal{L}$ is as in §7, that is,

$$\mathcal{L}(,\ -\,, \cdots, (\,+\,|\,-\,)/, \cdots, |\,-\,, |) = \,, \tag{13.12}$$

$$\mathcal{L}\,((\cdots, \backslash, \backslash +, \backslash, \cdots)) = \mathcal{L}\,((\cdots, \backslash, \backslash -, \backslash, \cdots)) + . \tag{13.13}$$

It is easy to see that $\Sigma^k(i,j) = \mathbb{C}\sigma_{\neg}(\beth, \gimel)$ for each $\binom{k}{ij} \in \mathcal{B}$ and that $\{\,\sigma_k(i,j) | \binom{k}{ij} \in \mathcal{B}\}$ is a linear basis of Σ. Since $\dim \Sigma^k(0,l) = \dim \Sigma^k(l,0) = \delta_{kl}$, we have $\Sigma^k \cong L_k \cong (\Sigma^k)^\vee$ by Theorem 7.5 and (3.7) . More explicitly, we have the following.

Proposition 13.1. *The map* $\Sigma^k \to (\Sigma^k)^\vee$; $\sigma_k(i,j) \mapsto c\binom{k}{ij}\sigma_k^\vee(i,j)$ *gives an identification of* $\mathfrak{S}(A_1;t)_\epsilon$*-comodules, where* $\{\sigma_k^\vee(j,i)\}$ *denotes the dual basis of* $\{\sigma_k(i,j)\}$ *and the constant* $c\binom{k}{ij}$ *is given by*

$$c\binom{k}{ij} = (-\epsilon)^{(i-j)/2}\frac{[(i+j+k)/2+1]!\,[(i-j+k)/2]!\,[(-i+j+k)/2]!}{[i+1]\,[(i+j-k)/2]!}. \tag{13.14}$$

Under this identification, the maps d_{Σ^k} *and* b_{Σ^k} *in* (3.8)-(3.9) *are given by*

$$d_{\Sigma^k}(i) = \sum_j c\binom{k}{ji}^{-1} \sigma_k(i,j) \otimes \sigma_k(j,i), \tag{13.15}$$

$$b_{\Sigma^k}\,(\sigma_k(i,j) \otimes \sigma_k(j,l)) = \delta_{il}\, c\binom{k}{ij}\, i \tag{13.16}$$

respectively, where the summation in (13.15) *is taken over all* $j \in \mathcal{V}$ *such that* $\binom{k}{ji} \in \mathcal{B}$.

Proof. It suffices to show the first assertion. Since $\Sigma^k(i,j) = \mathbb{C}\sigma_{\neg}(\beth, \gimel)$ and $(\Sigma^k)^\vee(i,j) = \mathbb{C}\sigma^\vee_{\neg}(\beth, \gimel)$, there exists an isomorphism $\Sigma^k \cong (\Sigma^k)^\vee$ of the form $\sigma_k(i,j) \mapsto c\binom{k}{ij}\sigma_k^\vee(i,j)$ for some nonzero constant $c\binom{k}{ij}$ ($\binom{k}{ij} \in \mathcal{B}$). To compute $c\binom{k}{ij}$, we consider the $\mathfrak{S}(A_1;t)_\epsilon$-right module structure on Σ given by (8.1). Similarly to Lemma 8.2, we obtain

$$\sigma_k(i,j)\, e\binom{i,\, i \pm 1}{j,\, j \pm 1} = \zeta^{-k}\epsilon^{(\pm i \mp j + k)/2} t^{(\mp i \pm j + k)/2} \frac{[\frac{i+j\mp k}{2} + 1]}{[i+1]}\sigma_k(i \pm 1, j \pm 1). \tag{13.17}$$

On the other hand, by (10.1), we obtain

$$S\left(e\binom{i,\, i \pm 1}{j,\, j \pm 1}\right) = \frac{[j+1]}{[i+1]}\, e\binom{j \pm 1,\, j}{i \pm 1,\, i}, \quad S\left(e\binom{i,\, i \pm 1}{j,\, j \mp 1}\right) = -\epsilon\, \frac{[j+1]}{[i+1]}\, e\binom{j \mp 1,\, j}{i \pm 1,\, i}. \tag{13.18}$$

Using this together with (8.3) and $(13.17)_-$, we obtain

$$\sigma_k^\vee(i,j)\, e\binom{i,i+1}{j,j+1} = \zeta^{-k}\epsilon^{(i-j+k)/2}t^{(-i+j+k)/2}\frac{[\frac{i+j+k}{2}+2]}{[i+2]}\sigma_k^\vee(i+1,j+1). \tag{13.19}$$

By $(13.17)_+$ and (13.19), we obtain

$$\frac{c\binom{k}{i+1,j+1}}{c\binom{k}{i,j}} = \frac{[i+1]\,[(i+j+k)/2+2]}{[i+2]\,[(i+j-k)/2+1]}. \tag{13.20}$$

Similarly, by computing $\sigma_k(i,j)\, e\binom{i,i\pm1}{j,j\mp1}$, we obtain

$$\frac{c\binom{k}{i\pm1,j\mp1}}{c\binom{k}{i,j}} = -\epsilon\,\frac{[i+1]\,[(\pm i\mp j+k)/2+1]}{[i+1\pm1]\,[(\mp i\pm j+k)/2]}. \tag{13.21}$$

By solving these recursion relations under some initial condition, we get (13.14). □

Proposition 13.2. (1) *The braiding* $c\colon \Sigma^m\bar{\otimes}\Sigma^n\tilde{\rightarrow}\Sigma^n\bar{\otimes}\Sigma^m$ *and its inverse are given by*

$$c^{\pm1}\left(\sigma_m(h,i)\otimes\sigma_n(i,k)\right) = \sum_j w^\pm_{mn}\begin{bmatrix} h & i \\ j & k\end{bmatrix}\sigma_n(h,j)\otimes\sigma_m(j,k), \tag{13.22}$$

$$w^\pm_{mn}\begin{bmatrix} h & i \\ j & k\end{bmatrix} := \sum_{\mathbf{u}\in\mathcal{G}_|^\setminus}\sum_{\mathbf{v}\in\mathcal{G}_|}\epsilon^{\mathcal{L}(\mathbf{u})+\mathcal{L}(\mathbf{v})+\mathcal{L}(\mathbf{q})+\mathcal{L}(\mathbf{r})}w^\pm\begin{bmatrix} & \mathbf{q} & \\ \mathbf{u} & & \mathbf{r} \\ & \mathbf{v} & \end{bmatrix}, \tag{13.23}$$

where $\mathbf{q}$ *and* $\mathbf{r}$ *denote arbitrary elements of* $\mathcal{G}$ *and* $\mathcal{G}^\setminus$ *respectively, and the summation in* (13.22) *is taken over all* $j\in\mathcal{V}$ *such that* $\binom{n}{hj},\binom{m}{jk}\in\mathcal{B}$.
(2) *The ribbon functional of* $\mathfrak{S}(\mathfrak{A}_1,\mathfrak{t})^\iota_{\epsilon,\zeta}$ *acts on* L_i *as the scalar* θ_i^{-1} *given by*

$$\theta_i = \iota^i\zeta^{i(i+2)}. \tag{13.24}$$

Proof. Since the braiding is a natural transformation and the map $\mathbb{C}\mathcal{G}\to\Sigma$; $\mathbf{p}\mapsto\sigma(\mathbf{p})$ is a $\mathfrak{S}(A_1;t)_\epsilon$-comodule map, we have

$$c^{\pm1}\left(\sigma_m(h,i)\otimes\sigma_n(i,k)\right) = \sum_j\sum_{\mathbf{u}\in\mathcal{G}_|^\setminus}\sum_{\mathbf{v}\in\mathcal{G}_|}\epsilon^{\mathcal{L}(\mathbf{q})+\mathcal{L}(\mathbf{r})}w^\pm\begin{bmatrix} & \mathbf{q} & \\ \mathbf{u} & & \mathbf{r} \\ & \mathbf{v} & \end{bmatrix}\sigma(\mathbf{u})\otimes\sigma(\mathbf{v}). \tag{13.25}$$

Since $\sigma(\mathbf{p}) = \epsilon^{\mathcal{L}(\mathbf{p})}\sigma_{\mathbf{m}}(\mathbf{i},\mathbf{j})$ for each $\mathbf{p}\in\mathcal{G}_|$, this proves Part (1). When $i=1$, (13.24) follows from the proof of Proposition 10.2. For $i>1$, (13.24) follows from (11.7) by induction on i. □

14. THE STATE SUM INVARIANTS

Let L be a positive integer and let ϵ,ι be elements of $\{\pm1\}$ such that $\epsilon=1$ if L is a odd integer. Let t be a primitive $2(L+2)$-th root of unity and ζ a solution of $\zeta^2=\epsilon\, t$. Applying the general theory of TQFT to the MTC $\mathcal{C}_{\mathfrak{S}}(\mathcal{A}_\infty,\sqcup)^\iota_{\epsilon,\zeta}$, we obtain an invariant $\tau^\iota_{\epsilon,\zeta}$ of oriented 3-manifolds. To give an explicit description of $\tau^\iota_{\epsilon,\zeta}$, we prepare some terminologies on link diagrams.

Let D be a generic link diagram in $\mathbb{R}\times(\nvdash,\nVdash)$ (viewed as a union of line segments $\overline{AB}$), which presents a framed link L with components $K_1,\dots,K_p$ (see e.g. [26]). A point of D is called *extremal* if the height function on D attains its local maximum or local minimum in this point, where the height function ht is the restriction of the projection $\mathbb{R}\times(\nvdash,\nVdash)\to(\nvdash,\nVdash)$ on D. A point of D is called *singular* if it is either an extremal point or a crossing point. We denote by $\sharp D$ the set of all singular points of D. Let $\mathcal{E}$ be the set of all connected components of $D\setminus\sharp D$. We say that $E\in\mathcal{E}$ *belongs* to K_q $(1\le q\le p)$ if E is a subset of the image of K_q via the projection $L \to D$. Let c be a map from $\{K_1, \dots, K_p\}$ to $\mathcal{V}$. We say that a map $\lambda \colon \mathcal{E} \to \mathcal{B}$; $E \mapsto \binom{\lambda_1(E)}{\lambda_2(E)\lambda_3(E)}$ is a *state* on D of *color c* if $\lambda_1(E) = c(K_q)$ for each component K_q and $E \in \mathcal{E}$ belonging to K_q. We denote by $\mathbf{c}_\lambda$ the color of a state λ, and by $\mathcal{S}(\mathcal{D})$ the set of all states on D. Figure (A) shows a state λ on a diagram of the Hopf link with 6 singular points, such that $\mathbf{c}_\lambda(\mathbf{K_1}) = \mathbf{1}$, $\mathbf{c}_\lambda(\mathbf{K_2}) = \mathbf{2}$.



Figure (A)

Next, we assign a complex number $\langle \lambda | A \rangle$ for each state $\lambda \in \mathcal{S}(\mathcal{D})$ and singular point $A \in \sharp D$ as follows: When (λ, A) is as in Figure (B) or Figure (C), then we set

$$\langle \lambda | A \rangle = c\binom{l}{hi}\delta_{hk}\delta_{ij} \quad \text{or} \quad \langle \lambda | \mathrm{A} \rangle = \mathrm{c}\binom{\mathrm{l}}{\mathrm{ih}}^{-1}\delta_{\mathrm{hk}}\delta_{\mathrm{ij}} \tag{14.1}$$

respectively, where $c\binom{l}{hi}$ is as in (13.14).



Figure (B)



Figure (C)



Figure (D_+)



Figure (D_-)

When (λ, A) is as in Figure ($D_\pm$), we set

$$\langle \lambda | A \rangle = \delta_{ij}\delta_{hc}\delta_{de}\delta_{fk} w^{\pm}_{mn}\begin{bmatrix} h & i \\ e & f \end{bmatrix}, \tag{14.2}$$

where $w^{\pm}_{mn}$ is as in (13.23).

The following result follows from §13 by a method quite similar to "vertex models on link invariants" (see e.g. [40] Appendix II), hence we omit the proof.

Theorem 14.1. *Let $\tau^{\iota}_{\epsilon,\zeta}$ be the invariant of closed oriented 3-manifolds associated with the modular tensor category $\mathcal{C}_{\mathfrak{S}}(\mathcal{A}_\infty, \sqcup)^{\iota}_{\epsilon,\zeta}$ (cf. [40]). Let M be a 3-manifold obtained by surgery on S^3 along a framed link L with p components $K_1, \ldots, K_p$. Let D be a generic diagram in $\mathbb{R} \times (\nvdash, \nVdash)$ which presents L. Then $\tau^{\iota}_{\epsilon,\zeta}$ is given by*

$$\tau^{\iota}_{\epsilon,\zeta}(M) = \Delta^{\sigma(L)}\mathcal{D}^{-\sigma(\mathcal{L})--}\sum_{\lambda\in\mathcal{S}(\mathcal{D})}\prod_{=}\iota^{\mathbf{c}_\lambda(\mathbf{K_q})}\left[\mathbf{c}_\lambda(\mathbf{K_q})+\mathbf{1}\right]\prod_{\mathbf{A}\in\sharp\mathbf{D}}\langle\lambda|\mathbf{A}\rangle. \tag{14.3}$$

Here Δ denotes a fixed square root of $\sum_{i\in\mathcal{V}}[i+1]^2$, $\mathcal{D} = \sum_{\in\mathcal{V}} \iota\zeta^{-(+)}[+]$ and $\sigma(L)$ denotes the signature of the linking matrix of L.

REFERENCES

[1] H. Andersen, Tensor products of quantized tilting modules, *Commun. Math. Phys.* **149** (1992), 149-159.
[2] G. Bergman, The diamond lemma for ring theory, *Adv. Math.* **29** (1978), 178-218.

[3] G. Böhm and K. Szlachányi, A coassociative C*-quantum group with non-integral dimensions, *Lett. Math. Phys.* **35** (1996), 437-456.

[4] V. Chari and A. Pressley, "A guide to quantum groups," Cambridge University Press, Cambridge, 1994.

[5] Y. Doi and M.Takeuchi, Multiplication alternation by two-cocycles — The quantum version —, *Commun. Alg.* **22** (1994), 5715-5732.

[6] V. G. Drinfeld, On quasitriangular Quasi-Hopf algebras and a group closely connected with $Gal(\bar{\mathbb{Q}}/\mathbb{Q})$ *Leningrad Math. J.* **2** (1991), 829-860.

[7] M. Finkelberg, An equivalence of fusion categories, *Geom. Funct. Analysis* **6** (1996), 249-267.

[8] S. Gelfand and D. Kazhdan, Examples of tensor categories, *Invent. Math.* **109** (1992), 595-617.

[9] F. Goodman and T. Nakanishi, Fusion algebras in integrable systems in two dimensions, *Phys. Lett. B* **262** (1991), 259-264.

[10] F. Goodman and H. Wenzl, Littlewood-Richardson coefficients for Hecke algebras at roots of unity, *Adv. Math.* **82** (1990), 244-265.

[11] T. Hayashi, An algebra related to the fusion rules of Wess-Zumino-Witten models, *Lett. Math Phys.* **22** (1991), 291-296.

[12] T. Hayashi, Quantum deformation of classical groups, *Publ. RIMS, Kyoto Univ.* **28** (1992), 57 - 81.

[13] T. Hayashi, Quantum groups and quantum determinants, *J. Algebra* **152** (1992), 146-165.

[14] T. Hayashi, Quantum group symmetry of partition functions of IRF models and its application to Jones' index theory, *Commun. Math. Phys.* **157** (1993), 331-345.

[15] T. Hayashi, Face algebras and their Drinfeld doubles, *in* "Proceedings of Symposia in Pure Mathematics," Vol 56, Part 2, American Mathematical Society, 1994.

[16] T. Hayashi, Face algebras I — A generalization of quantum group theory, to appear in *J. Math. Soc. Japan.*

[17] T. Hayashi, Compact quantum groups of face type, *Publ. RIMS, Kyoto Univ.* **32** (1996), 351 - 369.

[18] T. Hayashi, Galois quantum groups of II$_1$-subfactors, preprint.

[19] T. Hayashi, Face algebras II — Standard generator theorems, in preparation.

[20] T. Hayashi, Quantum groups and quantum semigroups, to appear in *J. Algebra.*

[21] T. Hayashi, in preparation.

[22] T. Hayashi, in preparation.

[23] M. Jimbo, T. Miwa and M. Okado, Solvable lattice models related to the vector representation of classical simple Lie algebras, *Commun. Math. Phys.* **116** (1988), 507-525.

[24] B. Jurčo and P. Schupp, AKS scheme for face and Calgero-Moser-Sutherland type models, preprint.

[25] V. Kac, "Infinite dimensional Lie algebras," 3rd ed., Cambridge Univ. press, 1990.

[26] C. Kassel, "Quantum groups," Springer-Verlag, New York, 1995.

[27] D. Kazhdan and H. Wenzl, Reconstructing monoidal categories, *Adv. in Soviet Math.* **16** (1993),111-136.

[28] A. Kirillov, Jr, On an inner product in modular tensor categories, *J. of AMS* **9** (1996),1135-1169.

[29] T. Koornwinder, Compact quantum groups and q-special functions, preprint.

[30] R. Larson and J. Towber, Two dual classes of bialgebras related to the concepts of "quantum group" and "quantum Lie algebra," *Commun. Alg.* **19** (1991), 3295-3345.

[31] I. Macdonald, "Symmetric functions and Hall polynomials," 2nd ed., Oxford Univ. press, Oxford, 1995.

[32] G. Moore and N. Seiberg, Classical and quantum conformal field theory, *Commun. Math. Phys.* **123** (1989), 177-254.

[33] A. Ocneanu, Quantized group, string algebras and Galois theory for algebras, In "Operator algebras and applications, Vol2". London Math. Soc. Lecture note series **136** (1989), 119-172.

[34] N. Reshetikhin and V. Turaev, Invariants of 3-manifolds via link polynomials and quantum groups, *Invent. Math.* **103** (1991), 547-598.

[35] N. Reshetikhin, L. Takhtadzhyan and L. Faddeev, Quantization of Lie groups and Lie algebras, *Leningrad Math. J.* **1** (1990), 193-225.

[36] P. Schauenburg, Face algebras are $\times_R$-bialgebras, preprint.

[37] M. Sweedler, "Hopf algebras," Benjamin Inc., New York, 1969.

[38] M. Takeuchi, Matric bialgebras and quantum groups, *Israel J. Math.* **72** (1990), 232-251.

[39] A. Tsuchiya and Y. Kanie, Vertex operators in conformal field theory on $\mathbb{P}^1$ and monodromy representations of braid groups, In "Adv. Stud. Pure Math. Vo16".

[40] V. Turaev, "Quantum invariants of knots and 3-manifolds," Walter de Gruyter, Berlin, New York, 1994.

[41] V. Turaev and H. Wenzl, Quantum invariants of 3-manifolds associated with classical simple Lie algebras, *Int. J. of Modern Math.* **4** (1993), 323-358.

[42] V. Turaev and H. Wenzl, Semisimple and modular categories from link invariants, *Math. Ann* **309** (1997), 411-461.

[43] E. Verlinde, Fusion rules and modular transformations in 2D conformal field theory, *Nucl. Phys.* **B300** (1988), 360-376.

[44] M. Walton, Algorithm for WZW fusion rules: A proof, *Phys. Lett. B* **241** (1990), 365-368.

[45] H. Wenzl, Hecke algebras of type A_n and subfactors, *Invent. Math.* **92** (1988), 349-383.

[46] H. Wenzl, C^* tensor categories from quantum groups, *J. AMS* **11** (1998), 261-282.

[47] E. Witten, Quantum field theory and the Jones polynomial, *Comm. Math. Phys.* **121** (1989), 351-399.